# Sensitivity analysis of the energy functional

JÚLIA SALAMON

ABSTRACT.

In the present work a numerical method is developed for the determination of the $J \in (0, \infty)$ interval in the three critical points theorem. The critical surfaces are also approximated.

REFERENCES

[1] Bonanno, G., *Some remarks on a three critical points theorem*, Nonlinear Analysis **54** (2003), 651 - 665
[2] Giaquinta, M. and Hildebrandt S., *Calculus of Variations I, The Lagrangian Formalism*, Springer, Berlin, 1996
[3] Hahn, B. D. and Valentine, D. T., *Essential matlab for engineers and scientists*, Elsevier, Oxford, 2007
[4] Kristály, A., *Multiplicity Results for an Eigenvalue Problem for Hemivariational Inequalities in Strip-Like Domains*, Set-Valued Analysis **13** (2005), 85 - 103
[5] Kristály, A., Existence of two non-trivial solutions for a class of quasilinear elliptic variational systems on strip-like domains, *Proceedings of the Edinburgh Mathematical Society* **48**, (2005) 1-13
[6] Kristály, A. and Varga, Cs., *On a class of quasilinear eigenvalue problems in RN*, Math. Nachr. **278**, No. 15, (2005) 1756 - 1765
[7] Ricceri, B., *On a three critical points theorem*, Arch. Math. **75** (2000), 220 - 226
[8] Tehrani, H., *Existence results for an indefinite unbounded perturbation of a resonant Schödinger equation*, J. Differential Equations **236**, (2007) 1 - 28
[9] http://www.mathworks.com/support/books

SAPIENTIA UNIVERSITY
DEPARTMENT OF MATHEMATICS AND INFORMATICS
LIBERTĂŢII 1, 530104 MIERCUREA CIUC, ROMANIA
*E-mail address*: salamonjulia@sapientia.siculorum.ro