Semifixed sets for multivalued φ -contractions

IOANA CAMELIA TIŞE

Abstract.

The purpose of this paper is to present some semifixed set theorems for multivalued φ -contractions. Our results generalize some recent theorems by F. S. De Blasi. As an application, existence and uniqueness for the solution of a set integral equation is obtained.

REFERENCES

- [1] De Blasi, F. S., Semifixed sets of maps in hyperspaces with application to set differential equations, Set-Valued Analysis, 14 (2006), 263-272
- [2] V. Lakshmikantham, T. Gnana Bhaskar, and Devi J. Vasundhara, Theory of Set Differential Equations in Metric Spaces, Cambridge Scientific Publishers, 2006
- [3] Petruşel, A., Petruşel, G. and Moţ, G., Topics in nonlinear analysis and applications to mathematical economics, House of the Book Science, Cluj-Napoca, 2007
- [4] Petruşel, A., (ϵ, φ) locally contractive multivalued mappings and applications, Studia Univ. Babeş-Bolyai, 36 (1991), 101-110
- [5] Rus, I. A., Generalized contractions and applications, Cluj Univ. Press, Cluj-Napoca, 2001

BABEŞ-BOLYAI UNIVERSITY DEPARTMENT OF APPLIED MATHEMATICS KOGĂLNICEANU 1 400084 CLUJ-NAPOCA, ROMANIA *E-mail address*: ticamelia@yahoo.com

Received: 12.10.2008; In revised form: 27.02.2009; Accepted: 21.05.2009. 2000 Mathematics Subject Classification. 47H10. Key words and phrases. Compact convex set, multivalued operator, semifixed set, Hukuhara's derivative, set integral equation.