

Minimal structures, m -open functions and bitopological spaces

TAKASHI NOIRI AND VALERIU POPA

ABSTRACT.

By using m -open functions from a topological space into an m -space, we establish the unified theory for several weak forms of open functions between bitopological spaces.

REFERENCES

- [1] Bose, S., *Semi-open sets, semi-continuity and semi-open mappings in bitopological spaces*, Bull. Calcutta Math. Soc. **73** (1981), 237-246
- [2] Jelić, M., *A decomposition of pairwise continuity*, J. Inst. Math. Comput. Sci. Math. Ser. **3** (1990), 25-29
- [3] Jelić, M., *Feebly p -continuous mappings*, Suppl. Rend. Circ. Mat. Palermo (2) **24** (1990), 387-395
- [4] Jelić, M., *On some mappings of bitopological spaces*, Suppl. Rend. Circ. Mat. Palermo (2) **29** (1992), 483-491
- [5] Kar, A. and Bhattacharyya, P., *Bitopological preopen sets, precontinuity and preopen mappings*, Indian J. Math. **34** (1992), 295-309
- [6] Khedr, F. H., Al-Areefi S. M. and Noiri, T., *Precontinuity and semi-precontinuity in bitopological spaces*, Indian J. Pure Appl. Math. **23** (1992), 625-633
- [7] Sampath, Kumar, S., *Pairwise α -open, α -closed and α -irresolute functions in bitopological spaces*, Bull. Inst. Math. Acad. Sinica **21** (1993), 59-72
- [8] Lipski, T., *Quasicontinuous multivalued maps in bitopological spaces*, Slupskie Prace Mat. Przyrodnicze Slupsk **7** (1988), 3-31
- [9] Maheshwari S. N. and Prasad, R., *Semiopen sets and semi continuous functions in bitopological spaces*, Math. Notae **26** (1977/78), 29-37
- [10] Maki, H., Rao K. C. and Nagoor Gani, A., *On generalizing semi-open and preopen sets*, Pure Appl. Math. Sci. **49** (1999), 17-29
- [11] Nasef A. A. and Noiri, T., *Feeble open sets and feeble continuity in bitopological spaces*, An. Univ. Timișoara Ser. Mat.-Inform. **36** (1998), 79-88
- [12] Palaniappan N. and Pious Missier, S., *δ -semi-open sets in bitopological spaces*, J. Indian Acad. Math. **25** (2003), 193-207
- [13] Palaniappan N. and Pious Missier, S., *δ -preopen sets in bitopological spaces*, J. Indian Acad. Math. **25** (2003), 287-295
- [14] Popa V. and Noiri, T., *On M -continuous functions*, Anal. Univ. "Dunarea de Jos" Galati, Ser. Mat. Fiz. Mec. Teor. (2) **18** (23) (2000), 31-41
- [15] Popa V. and Noiri, T., *On the definitions of some generalized forms of continuity under minimal conditions*, Mem. Fac. Sci. Kochi Univ. Ser. A Math. **22** (2001), 9-18
- [16] Popa V. and Noiri, T., *A unified theory of weak continuity for functions*, Rend. Circ. Mat. Palermo (2) **51** (2002), 439-464
- [17] Veličko, N. V., *H-closed topological spaces*, Amer. Math. Soc. Transl. (2) **78** (1968), 103-118

2949-1 SHIOKITA-CHO, HINAGU
YATSUSHIRO-SHI, KUMAMOTO-KEN
869-5142 JAPAN
E-mail address: t.noiri@nifty.com

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF BACĂU
600114 BACĂU, ROMANIA
E-mail address: vpopa@ub.ro

Received: 02.07.2007; In revised form: 30.01.2008.; Accepted:

2000 Mathematics Subject Classification. 54C10, 54E55.

Key words and phrases. m -structure, m_X -open, (i, j) - m -open, m -open, bitopological space.