On the 1**-homogeneous non-linear connections**

ESMAEIL AZIZPOUR, SEDDIGHE DARSARAEI and SEYED SAJJAD POURMORTAZAVI

Abstract.

In this paper, we show that for every semispray S on a vector bundle $(R \times TM, \pi, R \times M)$, there are several sequences of semisprays and correspondingly several nonlinear connections associated to it. It is important to derive conditions on S, which guarantee that a sequence of nonlinear connections associated to S is constant. We show that the homogeneity condition for S yields the result.

References

- Leon, M. De. and Rodrigues, P. R., Dynamical connections and non-autonomous Lagrangian systems, Ann. Fac. Sci. Toulouse Math., (5) 9 (1988), No. 2, 171–181
- [2] Miron, R. and Anastasiei, M., The Geometry of Lagrange Spaces: Theory and Applications, Kluwer Academic Publishers, 1994
- [3] Vondra, A., Sprays and homogeneous connections on $R \times TM$, Arch. Math. (Brno), 28 (1992), 163–173

DEPARTMENT OF PURE MATHEMATICS UNIVERSITY OF GUILAN, SCIENCES FACULTY OF MATHEMATICAL PO BOX 1914, RASHT, IRAN *E-mail address*: eazizpour@guilan.ac.ir

Received: 16.03.2013; In revised form: 16.10.2013; Accepted: 01.11.2013 2010 *Mathematics Subject Classification*. 53C05, 58A30. Key words and phrases. *Nonlinear connections, spray*.