

Computing the Wiener index of graphs on triples

M. GHORBANI

ABSTRACT.

Let S be a set of size $n \geq 8$ and V be the set of all subsets of S of size 3. Three types of intersection graphs $G_i(n)$, $i = 0, 1, 2$, can be defined with the vertex set V whose Wiener indices will be calculated in this paper.

REFERENCES

- [1] Darafsheh, M. R., *Computing the topological indices of some graphs*, Acta Appl. Math., **110** (2010), 1225–1235
- [2] Darafsheh, M. R., The Wiener, Szeged and PI index of the triangle graph, in *Proceedings of the 5th Asian Mathematical Conference*, Kuala Lumpur, Malaysia (Keong, L. S. and Kamarul, H., Eds.), 2009, Vol. I, 81–89
- [3] Dobrynin, A. A., Entringer, R., and Gutman, I., *Wiener index of trees: Theory and applications*, Acta Appl. Math., **66** (2001), 211–249
- [4] Dobrynin, A. A., Gutman, I., Klavzar, S. K., and Zigert, P., *Wiener index of hexagonal systems*, Acta Appl. Math., **72** (2002), 247–294
- [5] Gutman, I., Furtula, B., and Belic, J., *Notes on the hyper-Wiener index*, J. Serb. Chem. Soc., **68** (2003), 943–948
- [6] Gutman, I., Yeh, Y. N., Lee, S. L., and Chen, J. C., *Wiener number of dendrimers*, MATCH Commun. Math. Comput. Chem., **30** (1994), 103–115
- [7] Harary, F., *Graph theory*, Addison Wesley, Reading, 1968
- [8] Hosoya, H., *Topological index. A newly proposed quantity characterising the topological nature of structural isomer of saturated hydrocarbons*, Bull. Chem. Soc. Jpn., **4** (1971), 2332–2339
- [9] Mohammadyari, R. and Darafsheh, M. R., *Topological indices of the double odd graph $2O_k$* , Creative Math. & Inf., **20** (2011), 163–170
- [10] Prabhakara, Rao, N. and Laxmi Prasanna, A., *On the Wiener index of pentachains*, Appl. Math. Sci., **2** (2008), 2443–2457
- [11] Wiener, H., *Strustural determination of paraffin boiling points*, J. Am. Chem. Soc., **69** (1947), 17–20

DEPARTMENT OF MATHEMATICS AND TECHNOLOGY
MAZANDARAN UNIVERSITY OF SCIENCE
P.O. BOX 11111, BEHSHAHR, IRAN
E-mail address: m_ghorbani@iust.ac.ir