

Sequences interpolating some geometric inequalities

DORIN ANDRICA and DAN ȘTEFAN MARINESCU

ABSTRACT. Using the geometric dynamic of an iterative process (Theorem 2.1), we obtain refinements to some famous geometric inequalities in a triangle by constructing interpolating sequences.

1. INTRODUCTION

For the inequality $a \leq b$, where a, b are real numbers, an increasing sequence $(u_n)_{n \geq 0}$ with the property $a = u_0 \leq u_1 \leq \dots \leq u_n \leq \dots \leq b$ and $u_n \rightarrow b$, is called *increasing interpolating sequence*. Similarly, a decreasing sequence $(v_n)_{n \geq 0}$ satisfying $a \leq \dots \leq v_n \leq \dots \leq v_1 \leq v_0 = b$ and $v_n \rightarrow a$, is a *decreasing interpolating sequence* for the inequality $a \leq b$.

In this paper we will construct interpolating sequences for some geometric inequalities in a triangle due to Euler, Mitrinović, Weitzenböck, Gordon, Curry, Finsler-Hadwiger, Pólya-Szegő, and Chen.

2. AN USEFUL RESULT ABOUT ITERATIONS IN THE CIRCUMCIRCLE

Let ABC be a triangle with the angles A, B, C measured in radians, with the length-sides a, b, c , the circumradius R , the inradius r , the semiperimeter s , and the area K . For the fixed nonnegative real numbers x, y, z with $x + y + z = 1$, define recursively the sequences $(A_n)_{n \geq 0}, (B_n)_{n \geq 0}, (C_n)_{n \geq 0}$ by $A_{n+1} = xA_n + yB_n + zC_n, B_{n+1} = zA_n + xB_n + yC_n, A_{n+1} = yA_n + zB_n + xC_n, A_0 = A, B_0 = B, C_0 = C, n = 0, 1, \dots$. Note that $A_n, B_n, C_n > 0$ and $A_n + B_n + C_n = \pi, n = 0, 1, \dots$. Therefore, we can consider the triangle $A_n B_n C_n$ with the angles A_n, B_n, C_n and having the same circumcircle as triangle $ABC, n = 1, 2, \dots$. Denote this triangle by \mathcal{T}_n . Let $a_n, b_n, c_n, R, r_n, s_n, K_n$ be the length-sides, the circumradius, the inradius, the semiperimeter, and the area of the triangle \mathcal{T}_n , respectively.

Let us mention that such recursive systems describing some dynamic geometries are considered by S. Abbot [1], G. Z. Chang and P. J. Davis [6], R. J. Clarke [9], J. Ding, L. R. Hitt and X-M. Zhang [11], L. R. Hitt and X-M. Zhang [14], and D. Ismailescu and J. Jacobs [15].

The first result is contained in the following theorem.

Theorem 2.1. *With the above notations, if at most one of x, y, z is equal to 0, then the sequences $(A_n)_{n \geq 0}, (B_n)_{n \geq 0}, (C_n)_{n \geq 0}$ are convergent and*

$$(2.1) \quad \lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}.$$

Proof. It is easy to see that the following matrix relations hold

$$(2.2) \quad \begin{pmatrix} A_n \\ B_n \\ C_n \end{pmatrix} = U^n \begin{pmatrix} A \\ B \\ C \end{pmatrix},$$

Received: 23.11.2018. In revised form: 07.12.2018. Accepted: 14.12.2018

2010 *Mathematics Subject Classification.* 26D05, 26D07, 26D020.

Key words and phrases. *recursive system, Jensen inequality, geometric inequalities in a triangle.*

Corresponding author: Dorin Andrica; dandrica@math.ubbcluj.ro

where U is the circulant matrix given by

$$(2.3) \quad U = \begin{pmatrix} x & y & z \\ z & x & y \\ y & z & x \end{pmatrix}.$$

A simple induction argument shows that

$$(2.4) \quad U^n = \begin{pmatrix} x_n & y_n & z_n \\ z_n & x_n & y_n \\ y_n & z_n & x_n \end{pmatrix},$$

where the sequences $(x_n)_{n \geq 1}$, $(y_n)_{n \geq 1}$, $(z_n)_{n \geq 1}$ verify the recursive relations $x_{n+1} = xx_n + yy_n + zz_n$, $y_{n+1} = zx_n + xy_n + yz_n$, $z_{n+1} = yx_n + zy_n + xz_n$, $x_1 = x$, $y_1 = y$, $z_1 = z$, $n = 1, 2, \dots$. Summing down these relations we obtain $x_{n+1} + y_{n+1} + z_{n+1} = x_n + y_n + z_n$, hence the sequence $(x_n + y_n + z_n)_{n \geq 1}$ is constant and equal to 1.

On the other hand, the characteristic polynomial of the matrix U is

$$(2.5) \quad f_U(t) = (t - x - y - z)(t^2 + (y + z - 2x)t + x^2 + y^2 + z^2 - xy - yz - zx).$$

The hypothesis $x + y + z = 1$ implies that the roots of the polynomial f_U are $t_1 = 1$, $t_2 = \alpha$, $t_3 = \bar{\alpha}$, where $\alpha \in \mathbb{C} \setminus \mathbb{R}$ and $|\alpha| < 1$. It follows that we have

$$(2.6) \quad U = P \begin{pmatrix} 1 & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & \bar{\alpha} \end{pmatrix} P^{-1},$$

for some nonsingular matrix P . Therefore, we obtain

$$(2.7) \quad U^n = P \begin{pmatrix} 1 & 0 & 0 \\ 0 & \alpha^n & 0 \\ 0 & 0 & \bar{\alpha}^n \end{pmatrix} P^{-1},$$

and we get

$$x_n = a + b\alpha^n + c\bar{\alpha}^n, y_n = a' + b'\alpha^n + c'\bar{\alpha}^n, z_n = a'' + b''\alpha^n + c''\bar{\alpha}^n, n = 1, 2, \dots$$

for some fixed real numbers $a, b, c, a', b', c', a'', b'', c''$ determined by the initial conditions in the definition of the sequences $(x_n)_{n \geq 1}$, $(y_n)_{n \geq 1}$, $(z_n)_{n \geq 1}$. Because $\lim_{n \rightarrow \infty} \alpha^n = \lim_{n \rightarrow \infty} \bar{\alpha}^n = 0$, from the above formulas it follows that the sequences $(x_n)_{n \geq 1}$, $(y_n)_{n \geq 1}$, $(z_n)_{n \geq 1}$ are convergent and

$$\lim_{n \rightarrow \infty} x_n = a, \lim_{n \rightarrow \infty} y_n = a', \lim_{n \rightarrow \infty} z_n = a''.$$

From $x_n + y_n + z_n = 1$, $n = 1, 2, \dots$, we obtain $a + a' + a'' = 1$. On the other hand, the relation (2.7) shows that the eigenvalues of the matrix U^n are $1, \alpha^n, \bar{\alpha}^n$, that is the characteristic polynomial f_{U^n} of the matrix U^n is

$$f_{U^n}(t) = (t - x_n - y_n - z_n)(t^2 + (y_n + x_n - 2x_n)t + x_n^2 + y_n^2 + z_n^2 - x_n y_n - y_n z_n - z_n x_n)$$

and from the Vieta's relations we have $x_n^2 + y_n^2 + z_n^2 - x_n y_n - y_n z_n - z_n x_n = |\alpha|^{2n}$. When $n \rightarrow \infty$, we obtain the relation $a^2 + (a')^2 + (a'')^2 - aa' - a'a'' - a''a = 0$, i.e. $(a - a')^2 + (a' - a'')^2 + (a'' - a)^2 = 0$. Therefore $a = a' = a'' = \frac{1}{3}$, and the desired result follows from the relation (2). □

We will illustrate the above general iterative process by considering the following special geometric situation also studied by S. Abbot [1] and D. Șt. Marinescu, M. Monea, M. Opincariu and M. Stroe [18]. Recall that, if P is a point in the plane of the triangle ABC ,

the circumcevian triangle of P with respect to ABC is the triangle defined by the intersections of the Cevians AP, BP, CP with the circumcircle of ABC . We consider $A_1B_1C_1$ to be the circumcevian triangle of the incenter I of ABC , i.e. the circumcircle mid-arc triangle of ABC . In this case we have

$$A_1 = \frac{1}{2}(B + C), B_1 = \frac{1}{2}(C + A), C_1 = \frac{1}{2}(A + B),$$

that is in the general iterative process we have $x = 0, y = \frac{1}{2}, z = \frac{1}{2}$. On the other hand, because $A + B + C = \pi$, we have

$$A_1 = \frac{1}{2}(\pi - A), B_1 = \frac{1}{2}(\pi - B), C_1 = \frac{1}{2}(\pi - C)$$

Define recursively the sequence of triangles \mathcal{T}_n as follows : \mathcal{T}_{n+1} is the circumcircle mid-arc triangle with respect to the incenter of \mathcal{T}_n , and \mathcal{T}_0 is the triangle ABC . The angles of triangles \mathcal{T}_n are given by the recurrence relations $A_{n+1} = \frac{1}{2}(\pi - A_n), B_{n+1} = \frac{1}{2}(\pi - B_n), C_{n+1} = \frac{1}{2}(\pi - C_n)$, where $A_0 = A, B_0 = B, C_0 = C$. Solving these recurrences we get

$$\begin{aligned} A_n &= \left(-\frac{1}{2}\right)^n A + \frac{\pi}{3} \left(1 - \left(-\frac{1}{2}\right)^n\right), \\ B_n &= \left(-\frac{1}{2}\right)^n B + \frac{\pi}{3} \left(1 - \left(-\frac{1}{2}\right)^n\right), \\ C_n &= \left(-\frac{1}{2}\right)^n C + \frac{\pi}{3} \left(1 - \left(-\frac{1}{2}\right)^n\right), \end{aligned}$$

and the conclusion in Theorem 2.1 is obviously verified.

3. AN INTERPOLATING SEQUENCE FOR EULER'S INEQUALITY $R \geq 2r$

The Euler's inequality is a central result in triangle geometry (see T. Andreescu, O. Mushkarov and L. Stoyanov [3], D. Andrica [4], D. S. Mitrinovic, J. Pecaric and V. Volonec [20], and G. Popescu, I. V. Maftai, J. L. Diaz-Barrero and M. Dincă [22]). It is a direct consequence of Blundon's inequality and it has numerous and various refinements (see for instance D. Andrica [4], D. Andrica and D. Şt. Marinescu [5], and D. S. Mitrinovic, J. Pecaric and V. Volonec [20]). In this section we use the result in Theorem 2.1 to construct an increasing interpolating sequence for the Euler's inequality.

Theorem 3.2. With the above notations the sequence of inradii $(r_n)_{n \geq 0}$ is increasing and we have

$$(3.8) \quad \lim_{n \rightarrow \infty} r_n = \frac{R}{2}.$$

Proof. Using the known formula $\frac{r}{R} = 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$ we have to prove the following inequality

$$(3.9) \quad \sin \frac{A_{n+1}}{2} \sin \frac{B_{n+1}}{2} \sin \frac{C_{n+1}}{2} \geq \sin \frac{A_n}{2} \sin \frac{B_n}{2} \sin \frac{C_n}{2}.$$

Denote $\frac{A_n}{2} = u, \frac{B_n}{2} = v, \frac{C_n}{2} = t$ and the inequality (3.8) is equivalent to

$$(3.10) \quad \sin(xu + yv + zt) \sin(zu + xv + yt) \sin(yu + zv + xt) \geq \sin u \sin v \sin t.$$

To prove the inequality (3.10), let us consider the function $f : (0, \pi) \rightarrow \mathbb{R}$, defined by $f(s) = \ln \sin s$, which is concave on the interval $(0, \pi)$. Applying the Jensen's inequality we get the inequalities

$$\begin{aligned} f(xu + yv + zt) &\geq xf(u) + yf(v) + zf(t), \\ f(zu + xv + yt) &\geq zf(u) + xf(v) + yf(t), \end{aligned}$$

$$f(yu + zv + xt) \geq yf(u) + zf(v) + xf(t).$$

Summing these inequalities and using the relation $x + y + z = 1$, the inequality (3.10) follows.

From relation

$$\frac{r_n}{R} = 4 \sin \frac{A_n}{2} \sin \frac{B_n}{2} \sin \frac{C_n}{2}$$

and from (1) we obtain the limit (3.8). □

Corollary 3.1. With the above notations, the sequence of inradii $(r_n)_{n \geq 0}$ is an increasing interpolating sequence for the Euler's inequality, i.e. we have the inequalities

$$(3.11) \quad r = r_0 \leq r_1 \leq \dots \leq r_n \leq \dots \leq \frac{R}{2}.$$

4. AN INTERPOLATING SEQUENCE FOR MITRINOVIĆ'S INEQUALITY $s \leq \frac{3\sqrt{3}}{2}R$

The inequality $s \leq \frac{3\sqrt{3}}{2}R$ is known in the literature as Mitrinović's inequality. It is a simple consequence of Blundon's inequality but also there are different direct proofs. It has as a counterpart the inequality $3\sqrt{3}r \leq s$. Combining these two inequalities, we obtain a refinement to Euler's $R \geq 2r$:

$$3\sqrt{3}r \leq s \leq \frac{3\sqrt{3}}{2}R.$$

In what follows we use the result in Theorem 2.1 to construct an increasing interpolating sequence for the Mitrinović's inequality and a decreasing interpolating sequence for its counterpart.

Theorem 4.3. 1) With the above notations, the sequence of semiperimeters $(s_n)_{n \geq 0}$ is increasing and we have

$$(4.12) \quad \lim_{n \rightarrow \infty} s_n = \frac{3\sqrt{3}}{2}R.$$

2) The sequence $(\frac{s_n}{r_n})_{n \geq 0}$ is decreasing and

$$(4.13) \quad \lim_{n \rightarrow \infty} \frac{s_n}{r_n} = 3\sqrt{3}.$$

Proof. 1) We have $s_n = R(\sin A_n + \sin B_n + \sin C_n)$ and the function $g : (0, \pi) \rightarrow \mathbb{R}$, $g(u) = \sin u$ is concave on the interval $(0, \pi)$. From the Jensen's inequality we obtain $\sin A_{n+1} = \sin(xA_n + yB_n + zC_n) \geq x \sin A_n + y \sin B_n + z \sin C_n$. Similarly, we get other two inequalities $\sin B_{n+1} \geq z \sin A_n + x \sin B_n + y \sin C_n$ and $\sin C_{n+1} \geq y \sin A_n + z \sin B_n + x \sin C_n$. Summing up these inequalities it follows $s_n \leq s_{n+1}$. The relation $\lim_{n \rightarrow \infty} s_n = \frac{3\sqrt{3}}{2}R$ follows from Theorem 2.1.

2) From the relation $\cot \frac{A_n}{2} = \frac{s_n - a_n}{r_n}$ and the other two, we obtain

$$\cot \frac{A_n}{2} + \cot \frac{B_n}{2} + \cot \frac{C_n}{2} = \frac{3s_n - 2s_n}{r_n} = \frac{s_n}{r_n}.$$

Because the function $h(u) = \cot u$ is convex on the interval $(0, \frac{\pi}{2})$, with a similar argument as in the proof of part 1), it follows

$$\cot \frac{A_{n+1}}{2} + \cot \frac{B_{n+1}}{2} + \cot \frac{C_{n+1}}{2} \leq \cot \frac{A_n}{2} + \cot \frac{B_n}{2} + \cot \frac{C_n}{2},$$

that is $\frac{s_{n+1}}{r_{n+1}} \leq \frac{s_n}{r_n}$. The limit $\lim_{n \rightarrow \infty} \frac{s_n}{r_n} = 3\sqrt{3}$ follows from Theorem 2.1. □

Corollary 4.2. 1) With the above notations, the sequence of semiperimeters $(s_n)_{n \geq 0}$ is an increasing interpolating sequence for the Mitrinović's inequality, i.e. we have

$$(4.14) \quad s = s_0 \leq s_1 \leq \dots \leq s_n \leq \dots \leq \frac{3\sqrt{3}}{2}R.$$

2) The sequence $(\frac{s_n}{r_n})_{n \geq 0}$ is a decreasing interpolating sequence for the counterpart of Mitrinović's inequality, i.e. we have

$$(4.15) \quad 3\sqrt{3} \leq \dots \leq \frac{s_n}{r_n} \leq \dots \leq \frac{s_1}{r_1} \leq \frac{s_0}{r_0} = \frac{s}{r}.$$

5. INTERPOLATING WEITZENBÖCK'S INEQUALITY

In a triangle ABC , the Weitzenböck's inequality [25] is

$$(5.16) \quad a^2 + b^2 + c^2 \geq 4\sqrt{3}K,$$

where a, b, c are the length of the sides of the triangle and K denotes the area of ABC . To construct an interpolating sequence for (5.16) we use the special case $x = 0, y = \frac{1}{2}, y = \frac{1}{2}$ in the general iterative process described in Section 2. Considering the sequence $(u_n)_{n \geq 0}$, where

$$u_n = \frac{a_n^2 + b_n^2 + c_n^2}{4K_n}, n = 0, 1, \dots$$

we obtain the following result :

Theorem 5.4. With the above notations, the sequence $(u_n)_{n \geq 0}$ is decreasing and we have

$$(5.17) \quad \lim_{n \rightarrow \infty} u_n = \lim_{n \rightarrow \infty} \frac{a_n^2 + b_n^2 + c_n^2}{4K_n} = \sqrt{3}.$$

Proof. Clearly, we have

$$u_n = \frac{4R^2(\sin^2 A_n + \sin^2 B_n + \sin^2 C_n)}{8R^2 \sin A_n \sin B_n \sin C_n} = \frac{1}{2} \left(\frac{\sin A_n}{\sin B_n \sin C_n} + \frac{\sin B_n}{\sin C_n \sin A_n} + \frac{\sin C_n}{\sin A_n \sin B_n} \right) =$$

$$\frac{1}{2}(\cot B_n + \cot C_n + \cot C_n + \cot A_n + \cot A_n + \cot B_n) = \cot A_n + \cot B_n + \cot C_n.$$

It is easy to show that the inequality $2 \cot \frac{x+y}{2} \leq \cot x + \cot y$ holds for every $x, y \in (0, \pi)$ with $x + y < \pi$. Applying this property and using the recursive relations in the process, it follows

$$u_{n+1} = \cot A_{n+1} + \cot B_{n+1} + \cot C_{n+1} \leq \cot A_n + \cot B_n + \cot C_n = u_n,$$

that is the sequence is decreasing. Because $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}$, we obtain immediately the limit (5.17). \square

Corollary 5.3. With the above notations, the sequence $(u_n)_{n \geq 0}$ is a decreasing interpolating sequence for the Weitzenböck's inequality, i.e. we have

$$(5.18) \quad \sqrt{3} \leq \dots \leq u_n \leq \dots \leq u_1 \leq u_0 = \frac{a^2 + b^2 + c^2}{4K}.$$

6. INTERPOLATING GORDON'S INEQUALITY

In a triangle ABC the following inequality holds

$$(6.19) \quad ab + bc + ca \geq 4\sqrt{3}K,$$

where a, b, c are the length of the sides of the triangle and K denotes the area of ABC , and it is known as Gordon's inequality [12]. Denote by a_n, b_n, c_n, K_n the length of the sides and the area of the triangle $\mathcal{T}_n, n = 0, 1, 2, \dots$, where $a_0 = a, b_0 = b, c_0 = c$. Consider the sequence $(t_n)_{n \geq 0}$, where

$$t_n = \frac{a_n b_n + b_n c_n + c_n a_n}{K_n}, n = 0, 1, \dots$$

Theorem 6.5. With the above notations, the sequence $(t_n)_{n \geq 0}$ is decreasing and we have

$$(6.20) \quad \lim_{n \rightarrow \infty} t_n = \lim_{n \rightarrow \infty} \frac{a_n b_n + b_n c_n + c_n a_n}{K_n} = 4\sqrt{3}.$$

Proof. Using the area formula for triangle \mathcal{T}_n , we have

$$\frac{a_n b_n + b_n c_n + c_n a_n}{K_n} = 2 \left(\frac{1}{\sin A_n} + \frac{1}{\sin B_n} + \frac{1}{\sin C_n} \right).$$

Consequently, the property is equivalent to

$$\frac{1}{\sin A_n} + \frac{1}{\sin B_n} + \frac{1}{\sin C_n} \geq \frac{1}{\sin A_{n+1}} + \frac{1}{\sin B_{n+1}} + \frac{1}{\sin C_{n+1}}.$$

The function $f : (0, \pi) \rightarrow \mathbb{R}$, defined by $f(s) = \frac{1}{\sin s}$, is convex on the interval $(0, \pi)$. With a similar argument as in Theorem 3.2, the conclusion follows.

Because $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}$, we obtain the limit (6.20) from the first relation. \square

Corollary 6.4. With the above notations, the sequence $(t_n)_{n \geq 0}$ is an decreasing interpolating sequence for the Weitzenböck's inequality, i.e. we have

$$(6.21) \quad 4\sqrt{3} \leq \dots \leq t_n \leq \dots \leq t_1 \leq t_0 = \frac{ab + bc + ca}{K}.$$

7. INTERPOLATING CURRY'S INEQUALITY

Curry's inequality [10] is :

$$(7.22) \quad 4\sqrt{3}K \leq \frac{9abc}{a + b + c},$$

and it is an improvement to inequality (6.19).

Clearly, putting together these inequalities, we obtain the following interpolating inequalities to Weitzenböck inequality :

$$(7.23) \quad 4\sqrt{3}K \leq \frac{9abc}{a + b + c} \leq ab + bc + ca \leq a^2 + b^2 + c^2.$$

With the above notations, consider the sequence $(v_n)_{n \geq 0}$, where

$$v_n = \frac{9a_n b_n c_n}{4K_n (a_n + b_n + c_n)}, n = 0, 1, \dots$$

Theorem 7.6. The sequence $(v_n)_{n \geq 0}$ is decreasing and we have

$$(7.24) \quad \lim_{n \rightarrow \infty} v_n = \lim_{n \rightarrow \infty} \frac{9a_n b_n c_n}{4K_n(a_n + b_n + c_n)} = \sqrt{3}.$$

Proof. Because the relation

$$v_n = \frac{9R}{a_n + b_n + c_n} = \frac{9}{2(\sin A_n + \sin B_n + \sin C_n)},$$

and $\sin A_{n+1} = \sin(xA_n + yB_n + zC_n) \geq x \sin A_n + y \sin B_n + z \sin C_n$, $\sin B_{n+1} \geq z \sin A_n + x \sin B_n + y \sin C_n$, and $\sin C_{n+1} \geq y \sin A_n + z \sin B_n + x \sin C_n$, we get immediately the inequality $v_{n+1} \leq v_n$. The limit follows from Theorem 2.1. \square

Corollary 7.5. With the above notations, the sequence $(v_n)_{n \geq 0}$ is a decreasing interpolating sequence for the Curry's inequality, i.e. we have

$$(7.25) \quad \sqrt{3} \leq \dots \leq v_n \leq \dots \leq v_1 \leq v_0 = \frac{9abc}{4K(a+b+c)}.$$

8. INTERPOLATING FINSLER-HAWIGER INEQUALITY

It is well-known that in every triangle ABC the following inequality holds

$$(8.26) \quad a^2 + b^2 + c^2 \geq 4\sqrt{3}K + (a-b)^2 + (b-c)^2 + (c-a)^2$$

where a, b, c are the length of the sides of the triangle and K denotes the area of ABC . Clearly, it is a direct improvement to the Weitzenböck's inequality (5.16). The inequality (8.26) is known as the Finsler-Hawiger inequality [24], [13], and it is equivalent to

$$ab + bc + ca \geq 4\sqrt{3}K + \frac{1}{2}[(a-b)^2 + (b-c)^2 + (c-a)^2].$$

Therefore, it is also a strong improvement to the Gordon's inequality (6.19). The inequality (8.26) was intensively investigated by many authors ; C. Alsina and R. Nelsen [2], A. Cipu [9], C. Lupu and C. Pohoată [16], C. Lupu, C. Mateescu, V. Matei and M. Opincariu [17], and D. Șt. Marinescu, M. Monea, M. Opincariu and M. Stroe [19].

In what follows we use the result in Theorem 2.1 to construct a decreasing interpolating sequence for the Finsler-Hawiger inequality. With the above notations, consider the sequence $(w_n)_{n \geq 0}$, where

$$w_n = \frac{a_n^2 + b_n^2 + c_n^2 - (a_n - b_n)^2 - (b_n - c_n)^2 - (c_n - a_n)^2}{4K_n}, n = 0, 1, \dots$$

Theorem 8.7. The sequence $(w_n)_{n \geq 0}$ is decreasing and we have

$$(8.27) \quad \lim_{n \rightarrow \infty} w_n = \lim_{n \rightarrow \infty} \frac{a_n^2 + b_n^2 + c_n^2 - (a_n - b_n)^2 - (b_n - c_n)^2 - (c_n - a_n)^2}{4K_n} = \sqrt{3}.$$

Proof. Firstly, we will prove the following relation

$$(8.28) \quad \frac{a^2 + b^2 + c^2 - (a-b)^2 - (b-c)^2 - (c-a)^2}{4K} = \tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2}.$$

Indeed, starting with the left hand side, we have

$$\begin{aligned} \frac{2ab + 2bc + 2ca - a^2 - b^2 - c^2}{4K} &= \frac{1}{\sin A} + \frac{1}{\sin B} + \frac{1}{\sin C} - \frac{\sin A}{2 \sin B \sin C} - \frac{\sin B}{2 \sin C \sin A} \\ - \frac{\sin C}{2 \sin A \sin B} &= \frac{1}{\sin A} + \frac{1}{\sin B} + \frac{1}{\sin C} - \frac{1}{2}(\cot B + \cot C + \cot C + \cot A + \cot A + \cot B) = \end{aligned}$$

$$\frac{1}{\sin A} - \cot A + \frac{1}{\sin B} - \cot B + \frac{1}{\sin C} - \cot C = \tan \frac{A}{2} + \tan \frac{B}{2} + \tan \frac{C}{2},$$

since

$$\frac{1}{\sin A} - \cot A = \frac{1 - \cos A}{\sin A} = \frac{2 \sin^2 \frac{A}{2}}{2 \sin \frac{A}{2} \cos \frac{A}{2}} = \tan \frac{A}{2}.$$

Now, to prove that the sequence $(w_n)_{n \geq 0}$ is decreasing we are using relation (8.28) to obtain

$$w_{n+1} = \tan \frac{A_{n+1}}{2} + \tan \frac{B_{n+1}}{2} + \tan \frac{C_{n+1}}{2}.$$

The function $f : (0, \frac{\pi}{2}) \rightarrow \mathbb{R}$, defined by $f(s) = \tan s$, is convex on the interval $(0, \frac{\pi}{2})$. With a similar argument as in Theorem 2, we obtain the inequality $w_{n+1} \leq w_n$ and the conclusion follows.

Because $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}$ we obtain the limit (8.27) from the first relation. □

Corollary 8.6. With the above notations, the sequence $(w_n)_{n \geq 0}$ is a decreasing interpolating sequence for the Finsler-Hadwiger inequality, i.e. we have

$$(8.29) \quad \sqrt{3} \leq \dots \leq w_n \leq \dots \leq w_1 \leq w_0 = \frac{a^2 + b^2 + c^2 - (a-b)^2 - (b-c)^2 - (c-a)^2}{4K}.$$

9. INTERPOLATING PÓLYA-SZEGÖ INEQUALITY

Recall that in every triangle ABC the following inequality is due by Pólya and Szegő [21]:

$$(9.30) \quad K \leq \frac{\sqrt{3}}{4} (abc)^{\frac{2}{3}}.$$

Using our method we can construct an interpolating sequence for (21).

Theorem 9.8. With the above notations, the sequence $(\frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}})_{n \geq 0}$ is increasing and we have

$$(9.31) \quad \lim_{n \rightarrow \infty} \frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} = \frac{\sqrt{3}}{4}.$$

Proof. Using the area formula for triangle $A_n B_n C_n$, we have

$$\frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} = \frac{(a_n b_n c_n)^{\frac{1}{3}}}{4R_n} = \frac{\sqrt[3]{\sin A_n \sin B_n \sin C_n}}{2},$$

and the property is equivalent to

$$\sin A_n \sin B_n \sin C_n \leq \sin A_{n+1} \sin B_{n+1} \sin C_{n+1}.$$

From the argument in the proof of Theorem 3.2 for the concave function $f : (0, \pi) \rightarrow \mathbb{R}$ defined by $f(s) = \ln \sin s$, the conclusion follows. The limit (16) follows from the first relation and the limits $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}$. □

Corollary 9.7. With the above notations, the sequence $(\frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}})_{n \geq 0}$ is an increasing interpolating sequence for the Pólya-Szegő inequality, i.e. we have

$$(9.32) \quad \frac{K}{(abc)^{\frac{2}{3}}} \leq \frac{K_1}{(a_1 b_1 c_1)^{\frac{2}{3}}} \leq \dots \leq \frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \leq \dots \leq \frac{\sqrt{3}}{4}.$$

10. INTERPOLATING CHEN'S INEQUALITY

In the recent paper of Y-D. Wu, V. Lokesha and H. M. Srivastava [26] it is presented a refinement of inequality (5.16) in the form

$$(10.33) \quad K \leq \frac{\sqrt{3}}{4} (abc)^{\frac{2}{3}} \left(\frac{2r}{R}\right)^{\frac{1}{3}},$$

mentioning that this inequality is due by S.-L. Chen [7]. Using our method, we can obtain an interpolating sequence to this result.

Theorem 10.9. With the above notations, the sequence $\left(\frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \left(\frac{2r_n}{R_n}\right)^{\frac{1}{3}}\right)_{n \geq 0}$ is increasing and we have

$$(10.34) \quad \lim_{n \rightarrow \infty} \frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \left(\frac{2r_n}{R_n}\right)^{\frac{1}{3}} = \frac{\sqrt{3}}{4}.$$

Proof. The property does not result from the monotony of the two sequences because they are of the opposite monotony. Observe that

$$\begin{aligned} \frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \left(\frac{2r_n}{R_n}\right)^{\frac{1}{3}} &= \frac{\sqrt[3]{\sin A_n \sin B_n \sin C_n}}{2} \cdot \frac{1}{\sqrt[3]{8 \sin \frac{A_n}{2} \sin \frac{B_n}{2} \sin \frac{C_n}{2}}} = \\ &= \frac{\sqrt[3]{\cos \frac{A_n}{2} \cos \frac{B_n}{2} \cos \frac{C_n}{2}}}{2}. \end{aligned}$$

The function $f : (0, \pi/2) \rightarrow \mathbb{R}$ defined by $f(s) = \ln \cos s$ is concave, and we obtain as in the proof of Theorem 2, the inequality

$$\ln \cos \frac{A_{n+1}}{2} + \ln \cos \frac{B_{n+1}}{2} + \ln \cos \frac{C_{n+1}}{2} \geq \ln \cos \frac{A_n}{2} + \ln \cos \frac{B_n}{2} + \ln \cos \frac{C_n}{2}.$$

The limit (10.34) can be obtained by using the relations $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \lim_{n \rightarrow \infty} C_n = \frac{\pi}{3}$. \square

Corollary 10.8. The sequence $\left(\frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \left(\frac{2r_n}{R_n}\right)^{\frac{1}{3}}\right)_{n \geq 0}$ is an increasing interpolating sequence for the Chen's inequality, i.e. we have

$$(10.35) \quad \frac{K}{(abc)^{\frac{2}{3}}} \left(\frac{2r}{R}\right)^{\frac{1}{3}} \leq \frac{K_1}{(a_1 b_1 c_1)^{\frac{2}{3}}} \left(\frac{2r_1}{R_1}\right)^{\frac{1}{3}} \leq \dots \leq \frac{K_n}{(a_n b_n c_n)^{\frac{2}{3}}} \left(\frac{2r_n}{R_n}\right)^{\frac{1}{3}} \leq \dots \leq \frac{\sqrt{3}}{4}.$$

REFERENCES

- [1] Abbot, S., *Average sequences and triangles*, Math. Gaz., **80** (1996), 222–224
- [2] Alsina, C. and Nelsen, R., *Geometric proofs of the Weitzenböck and Finsler-Hadwiger inequality*, Math. Mag., **81** (2008), 216–219
- [3] Andreescu, T. Mushkarov, O. and Stoyanov, L., *Geometric Problems on Maxima and Minima*, Birkauer, Boston-Basel-Berlin, 2006
- [4] Andrica, D., *GEOMETRIE. Teme pentru perfecționarea profesorilor de matematică* (Romanian), Casa Cărții de Știință, Cluj-Napoca, 2017
- [5] Andrica, D. and Marinescu, D. Șt., *New Interpolation Inequalities to Euler's $R \geq 2r$* , Forum Geometricorum, Volume **17** (2017), 149–156
- [6] Chang, G. Z. and Davis, P. J., *Iterative processes in elementary geometry*, Amer. Math. Monthly, **90** (1983), No. 7, 421–431
- [7] Chen, S.-L., *An inequality chain relating to several famous inequalities*, Hunan Math. Comm., **1** (1995), 41 (in Chinese)
- [8] Cipu, A., *Optimal reverse Finsler-Hadwiger inequalities*, Gaz. Mat. Ser. A, **30** (2010), Nr. 3-4 (2012), 61–68
- [9] Clarke, R. J., *Sequences of polygons*, Math. Mag., **90** (1979), No. 2, 102–105

- [10] Curry, T. R., *Problem E 1861*, The American Mathematical Monthly, **73** (1966), No. 2
- [11] Ding, J., Hitt, L. R. and Zhang, X-M., *Markov chains and dynamic geometry of polygons*, Linear Algebra and Its Applications, **367** (2003), 255–270
- [12] Gordon, V. O., *Matematika v Skole*, 1966, No. 1, 89
- [13] Hadwiger, H., *Jber. Deutsch. Math. Verein.*, **49** (1939), 35–39
- [14] Hitt, L. R. and Zhang, X-M., *Dynamic geometry of polygons*, Elem. Math., **56** (2001), No. 1, 21–37
- [15] Ismailescu, D. and Jacobs, J., *On sequences of nested triangles*, Period. Math. Hungar., **53** (2006), No. 1-2, 169–184
- [16] Lupu, C. and Pohoățã, C., *Sharpening the Finsler-Hadwiger inequality*, Crux Mathematicorum with Mathematical Mayhem, **34** (2008), Issue 2, 97–101
- [17] Lupu, C., Mateescu, C., Matei, V. and Opincariu, M., *A refinement of the Finsler-Hadwiger reverse inequality*, Gaz. Mat. Ser. A, **28** (2010), 130–133
- [18] Marinescu, D. Șt., Monea, M., Opincariu, M. and Stroe, M., *A Sequence of Triangles and Geometric Inequalities*, Forum Geom., **9** (2009), 291–295
- [19] Marinescu, D. Șt., Monea, M., Opincariu, M. and Stroe, M., *Note on Hawiger-Finsler’s inequalities*, Math. Inequal. Appl., **6** (2012), No. 1, 57–64
- [20] Mitrinović, D. S., Pecarić, J. and Volonec, V., *Recent Advances in Geometric Inequalities (Mathematics and its Applications)*, Springer; Softcover reprint of the original 1st ed. 1989 edition (September 17, 2011)
- [21] Pólya, G. and Szegő, G., *Problems and Theorems in Analysis*, Vol. II, in: Grundlehren der Mathematischen Wissenschaften, Band 20, Springer-Verlag, New York, Heidelberg and Berlin, 1976, (Translated from the revised and enlarged Fourth German Edition by C. E. Billigheimer)
- [22] Popescu, P. G., Maftai, I. V., Diaz-Barrero, J. L. and Dincă, M., *Inegalități matematice: modele inovatoare* (Romanian), Editura Didactică și Pedagogică, București, 2007
- [23] Stoica, E., Minculete, N. and Barbu, C., *New aspects of Weitzenböck’s inequality*, Balkan J. Geom. Appl., **21** (2016), No. 2, pp. 95–101
- [24] Von Finsler, P. and Hadwiger, H., *Einige Relationen im Dreieck*, Commentarii Mathematici Helvetici, **10** (1937), No. 1, 316–326
- [25] Weitzenböck, R., *Über eine Ungleichung in der Dreiecksgeometrie*, Mathematische Zeitschrift, **5** (1919), No. 1–2, 137–146
- [26] Wu, Y-D., Loksha, V. and Srivastava, H. M., *Another refinement of the Pólya- Szegő inequality*, Computers and Mathematics with Applications, **60** (2010), 761–770

“BABEȘ-BOLYAI” UNIVERSITY
 FACULTY OF MATHEMATICS AND COMPUTER SCIENCE
 CLUJ-NAPOCA, ROMANIA
 Email address: dandrica@math.ubbcluj.ro

“IANCU DE HUNEDOARA” NATIONAL COLLEGE
 HUNEDOARA, ROMANIA
 Email address: marinescuds@gmail.com