

# Inequalities for the finite Hilbert transform of functions with bounded divided differences

SILVESTRU SEVER DRAGOMIR

**ABSTRACT.** In this paper we establish some inequalities for the finite Hilbert transform of complex valued functions for which the divided differences in any two points of the interval are bounded. Applications for some particular functions of interest are provided as well.

## 1. INTRODUCTION

Allover this paper, we consider the *finite Hilbert transform* on the open interval  $(a, b)$  defined by

$$(Tf)(a, b; t) := \frac{1}{\pi} PV \int_a^b \frac{f(\tau)}{\tau - t} d\tau := \lim_{\varepsilon \rightarrow 0^+} \left[ \int_a^{t-\varepsilon} + \int_{t+\varepsilon}^b \right] \frac{f(\tau)}{\pi(\tau - t)} d\tau$$

for  $t \in (a, b)$  and for various classes of functions  $f$  for which the above Cauchy Principal Value integral exists, see [13, Section 3.2] or [17, Lemma II.1.1].

For several recent papers devoted to inequalities for the finite Hilbert transform  $(Tf)$ , see [2]-[10], [14]-[16] and [18]-[19].

The following result holds.

**Theorem 1.1** (Dragomir et al., 2001 [1]). *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a monotonic nondecreasing (nonincreasing) function on  $[a, b]$ . If the finite Hilbert transform  $(Tf)(a, b, \cdot)$  exists in every  $t \in (a, b)$ , then*

$$(Tf)(a, b; t) \geq (\leq) \frac{1}{\pi} f(t) \ln \left( \frac{b-t}{t-a} \right) \quad (1.1)$$

for all  $t \in (a, b)$ .

The following result can be useful in practice.

**Corollary 1.1.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  and  $e : [a, b] \rightarrow \mathbb{R}$ ,  $e(t) = t$  such that  $f - me$ ,  $Me - f$  are monotonic nondecreasing, where  $m < M$  are given real numbers. If  $(Tf)(a, b, \cdot)$  exists in every point  $t \in (a, b)$ , then we have the inequality*

$$\frac{(b-a)m}{\pi} \leq (Tf)(a, b; t) - \frac{1}{\pi} f(t) \ln \left( \frac{b-t}{t-a} \right) \leq \frac{(b-a)M}{\pi} \quad (1.2)$$

for all  $t \in (a, b)$ .

**Remark 1.1.** If the function  $f$  is differentiable on  $(a, b)$  the condition that  $f - me$ ,  $Me - f$  are monotonic nondecreasing is equivalent with the following more practical condition

$$m \leq f'(t) \leq M \quad \text{for all } t \in (a, b). \quad (1.3)$$

---

Received: 21.03.2019. In revised form: 16.12.2019. Accepted: 10.01.2020

2010 *Mathematics Subject Classification.* 26D15, 26D10.

Key words and phrases. *finite Hilbert transform, divided difference, integral inequalities.*

From (1.2) we may deduce the following approximation result

$$\left| (Tf)(a, b; t) - \frac{1}{\pi} f(t) \ln \left( \frac{b-t}{t-a} \right) - \frac{M+m}{2\pi} (b-a) \right| \leq \frac{M-m}{2\pi} (b-a) \quad (1.4)$$

for all  $t \in (a, b)$ .

Motivated by the above results, in this paper we establish some inequalities for the finite Hilbert transform of complex valued functions for which the divided differences in any two points of the interval are bounded. Applications for some particular functions of interest are provided as well.

## 2. MAIN RESULTS

For a function  $f : (a, b) \rightarrow \mathbb{C}$  we define the *divided difference*

$$[f; t, s] := \frac{f(t) - f(s)}{t - s} \text{ for } t, s \in (a, b), t \neq s.$$

Now, for  $\gamma, \Gamma \in \mathbb{C}$  and  $(a, b)$  an interval of real numbers, define the sets of complex-valued functions

$$\bar{U}_{(a,b),d}(\gamma, \Gamma) := \left\{ f : (a, b) \rightarrow \mathbb{C} \mid \operatorname{Re} \left[ (\Gamma - [f; t, s]) \left( \overline{[f; t, s]} - \bar{\gamma} \right) \right] \geq 0, \right. \\ \left. \text{for all } t, s \in (a, b), t \neq s \right\} \quad (2.5)$$

and

$$\bar{\Delta}_{(a,b),d}(\gamma, \Gamma) := \left\{ f : (a, b) \rightarrow \mathbb{C} \mid \left| [f; t, s] - \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2} |\Gamma - \gamma| \right. \\ \left. \text{for all } t, s \in (a, b), t \neq s \right\}. \quad (2.6)$$

The following representation result may be stated.

**Proposition 2.1.** *For any  $\gamma, \Gamma \in \mathbb{C}$ ,  $\gamma \neq \Gamma$ , we have that  $\bar{U}_{(a,b),d}(\gamma, \Gamma)$  and  $\bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$  are nonempty, convex and closed sets and*

$$\bar{U}_{(a,b),d}(\gamma, \Gamma) = \bar{\Delta}_{(a,b),d}(\gamma, \Gamma). \quad (2.7)$$

*Proof.* We observe that for any  $z \in \mathbb{C}$  we have the equivalence

$$\left| z - \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2} |\Gamma - \gamma|$$

if and only if

$$\operatorname{Re} [(\Gamma - z)(\bar{z} - \bar{\gamma})] \geq 0.$$

This follows by the equality

$$\frac{1}{4} |\Gamma - \gamma|^2 - \left| z - \frac{\gamma + \Gamma}{2} \right|^2 = \operatorname{Re} [(\Gamma - z)(\bar{z} - \bar{\gamma})]$$

that holds for any  $z \in \mathbb{C}$ .

The equality (2.7) is thus a simple consequence of this fact.  $\square$

On making use of the complex numbers field properties we can also state that:

**Corollary 2.2.** *For any  $\gamma, \Gamma \in \mathbb{C}$ ,  $\gamma \neq \Gamma$ , we have that*

$$\bar{U}_{(a,b),d}(\gamma, \Gamma) = \left\{ f : (a, b) \rightarrow \mathbb{C} \mid (\operatorname{Re} \Gamma - \operatorname{Re} [f; t, s]) (\operatorname{Re} [f; t, s] - \operatorname{Re} \gamma) \right. \\ \left. + (\operatorname{Im} \Gamma - \operatorname{Im} [f; t, s]) (\operatorname{Im} [f; t, s] - \operatorname{Im} \gamma) \geq 0 \text{ for all } t, s \in (a, b), t \neq s \right\}. \quad (2.8)$$

Now, if we assume that  $Re(\Gamma) \geq Re(\gamma)$  and  $Im(\Gamma) \geq Im(\gamma)$ , then we can define the following set of functions as well:

$$\begin{aligned} \bar{S}_{(a,b),d}(\gamma, \Gamma) := \{f : (a, b) \rightarrow \mathbb{C} \mid Re(\Gamma) \geq Re[f; t, s] \geq Re(\gamma) \\ \text{and } Im(\Gamma) \geq Im[f; t, s] \geq Im(\gamma) \text{ for all } t, s \in (a, b), t \neq s\}. \end{aligned} \quad (2.9)$$

One can easily observe that  $\bar{S}_{(a,b)}(\gamma, \Gamma)$  is closed, convex and

$$\emptyset \neq \bar{S}_{(a,b),d}(\gamma, \Gamma) \subseteq \bar{U}_{(a,b),d}(\gamma, \Gamma). \quad (2.10)$$

The following result holds:

**Theorem 2.2.** *Let  $f : (a, b) \rightarrow \mathbb{C}$  be such that for some  $\gamma, \Gamma \in \mathbb{C}$ ,  $\gamma \neq \Gamma$ , we have that  $f \in \bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$ . Then we have the inequality*

$$\left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left( \frac{b-t}{t-a} \right) - \frac{1}{\pi} \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2\pi} |\Gamma - \gamma| \quad (2.11)$$

for any  $t \in (a, b)$ .

In particular, for  $t = \frac{a+b}{2}$  we obtain

$$\left| (Tf) \left( a, b; \frac{a+b}{2} \right) - \frac{1}{\pi} \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2\pi} |\Gamma - \gamma|. \quad (2.12)$$

*Proof.* Since  $f \in \bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$  it follows that

$$\left| f(t) - f(s) - \frac{\gamma + \Gamma}{2} (t - s) \right| \leq \frac{1}{2} |\Gamma - \gamma| |t - s|$$

for any  $t, s \in (a, b)$ .

By the continuity of the modulus property, we have

$$|f(t) - f(s)| - \left| \frac{\gamma + \Gamma}{2} \right| |t - s| \leq \left| f(t) - f(s) - \frac{\gamma + \Gamma}{2} (t - s) \right| \leq \frac{1}{2} |\Gamma - \gamma| |t - s|,$$

for any  $t, s \in (a, b)$ , which implies that

$$|f(t) - f(s)| \leq \frac{1}{2} (|\gamma + \Gamma| + |\Gamma - \gamma|) |t - s|$$

for any  $t, s \in (a, b)$ , showing that  $f$  is also Lipschitzian on  $(a, b)$ . Therefore, we conclude that the finite Hilbert transform  $T(f)(a, b; t)$  exists for all  $t \in (a, b)$ , see [13, Section 3.2] or [17, Lemma II.1.1].

For the mapping,  $\mathbf{1}(t) = 1$ ,  $t \in (a, b)$ , we have

$$\begin{aligned} (T\mathbf{1})(a, b; t) &= \frac{1}{\pi} PV \int_a^b \frac{1}{\tau - t} d\tau \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0^+} \left[ \int_a^{t-\varepsilon} \frac{1}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{1}{\tau - t} d\tau \right] \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0^+} \left[ \ln |\tau - t| \Big|_a^{t-\varepsilon} + \ln(\tau - t) \Big|_{t+\varepsilon}^b \right] \\ &= \frac{1}{\pi} \lim_{\varepsilon \rightarrow 0^+} [\ln \varepsilon - \ln(t - a) + \ln(b - t) - \ln \varepsilon] \\ &= \frac{1}{\pi} \ln \left( \frac{b - t}{t - a} \right), \quad t \in (a, b). \end{aligned}$$

Then, obviously, for  $f : (a, b) \rightarrow \mathbb{R}$  we have

$$\begin{aligned} (Tf)(a, b; t) &= \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t) + f(t)}{\tau - t} d\tau \\ &= \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau + \frac{f(t)}{\pi} PV \int_a^b \frac{1}{\tau - t} d\tau \end{aligned}$$

from where we get the equality

$$(Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left( \frac{b-t}{t-a} \right) = \frac{1}{\pi} PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \quad (2.13)$$

for any  $t \in (a, b)$ .

Since  $f \in \bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$ , hence

$$\begin{aligned} &\left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left( \frac{b-t}{t-a} \right) - \frac{1}{\pi} \frac{\gamma + \Gamma}{2} \right| \\ &= \left| \frac{1}{\pi} PV \int_a^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right) d\tau \right| \\ &\leq \frac{1}{\pi} PV \int_a^b \left| \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right| d\tau \leq \frac{1}{2} |\Gamma - \gamma| \frac{1}{\pi} PV \int_a^b d\tau \\ &= \frac{1}{2\pi} |\Gamma - \gamma| \end{aligned}$$

and the inequality (2.11) is thus obtained.  $\square$

**Remark 2.2.** We observe that if  $f - me$ ,  $Me - f$  are monotonic nondecreasing, where  $m < M$  are given real numbers, then we have that  $f \in \bar{\Delta}_{(a,b),d}(m, M)$  and from (2.11) we recapture (1.2).

We need the following technical lemma:

**Lemma 2.1.** Let  $f : (a, b) \rightarrow \mathbb{C}$  and  $t \in (a, b)$ . Provided that all integrals below exists, we have for any  $\delta \in \mathbb{C}$  that

$$\begin{aligned} &\int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \\ &+ 2 \left( \frac{1}{t+\varepsilon-a} \int_a^{t-\varepsilon} f(\tau) d\tau - \frac{1}{b-t+\varepsilon} \int_{t+\varepsilon}^b f(\tau) d\tau \right) \\ &+ 2 \left( \frac{b-t-\varepsilon}{b-t+\varepsilon} - \frac{t-\varepsilon-a}{t+\varepsilon-a} \right) f(t) \\ &= \frac{2}{t+\varepsilon-a} \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{a+t-\varepsilon}{2} \right) d\tau \\ &- \frac{2}{b-t+\varepsilon} \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{b+t+\varepsilon}{2} \right) d\tau, \end{aligned} \quad (2.14)$$

where  $\varepsilon > 0$  and such that  $\min \{t-a, b-t\} > \varepsilon$ .

*Proof.* We have for any  $\delta \in \mathbb{C}$  that

$$\begin{aligned}
 & \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} (\tau - t) d\tau \\
 & - \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau \frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} (\tau - t) d\tau \\
 & = \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - t - \frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} (s - t) ds \right) d\tau \\
 & = \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{a + t - \varepsilon}{2} \right) d\tau
 \end{aligned} \tag{2.15}$$

for  $t - a > \varepsilon > 0$ .

Since

$$\int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} (\tau - t) d\tau = \int_a^{t-\varepsilon} f(\tau) d\tau - (t - \varepsilon - a) f(t)$$

and

$$\frac{1}{t - \varepsilon - a} \int_a^{t-\varepsilon} (\tau - t) d\tau = -\frac{t + \varepsilon - a}{2},$$

then by (2.15) we get

$$\begin{aligned}
 & \int_a^{t-\varepsilon} f(\tau) d\tau - (t - \varepsilon - a) f(t) + \frac{t + \varepsilon - a}{2} \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau \\
 & = \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{a + t - \varepsilon}{2} \right) d\tau
 \end{aligned}$$

from where we obtain

$$\begin{aligned}
 & \frac{2}{t + \varepsilon - a} \int_a^{t-\varepsilon} f(\tau) d\tau - 2 \left( \frac{t - \varepsilon - a}{t + \varepsilon - a} \right) f(t) + \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau \\
 & = \frac{2}{t + \varepsilon - a} \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{a + t - \varepsilon}{2} \right) d\tau,
 \end{aligned}$$

namely

$$\begin{aligned}
 & \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau + \frac{2}{t + \varepsilon - a} \int_a^{t-\varepsilon} f(\tau) d\tau - 2 \left( \frac{t - \varepsilon - a}{t + \varepsilon - a} \right) f(t) \\
 & = \frac{2}{t + \varepsilon - a} \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{a + t - \varepsilon}{2} \right) d\tau,
 \end{aligned} \tag{2.16}$$

for  $t - a > \varepsilon > 0$ .

We have for any  $\delta \in \mathbb{C}$  that

$$\begin{aligned}
 & \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} (\tau - t) d\tau - \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \frac{1}{b - \varepsilon - t} \int_{t+\varepsilon}^b (\tau - t) d\tau \\
 & = \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - t - \frac{1}{b - \varepsilon - t} \int_{t+\varepsilon}^b (s - t) ds \right) d\tau \\
 & = \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{b + t + \varepsilon}{2} \right) d\tau
 \end{aligned} \tag{2.17}$$

for  $b - t > \varepsilon > 0$ .

Since

$$\int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} (\tau - t) d\tau = \int_{t+\varepsilon}^b f(\tau) d\tau - (b - t - \varepsilon) f(t)$$

and

$$\frac{1}{b - \varepsilon - t} \int_{t+\varepsilon}^b (\tau - t) d\tau = \frac{b - t + \varepsilon}{2},$$

then by (2.17) we get

$$\begin{aligned} & \int_{t+\varepsilon}^b f(\tau) d\tau - (b - t - \varepsilon) f(t) - \frac{b - t + \varepsilon}{2} \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \\ &= \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - t - \frac{1}{b - \varepsilon - t} \int_{t+\varepsilon}^b (s - t) ds \right) d\tau \\ &= \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{b + t + \varepsilon}{2} \right) d\tau, \end{aligned}$$

namely

$$\begin{aligned} & \frac{2}{b - t + \varepsilon} \int_{t+\varepsilon}^b f(\tau) d\tau - 2 \left( \frac{b - t - \varepsilon}{b - t + \varepsilon} \right) f(t) - \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \\ &= \frac{2}{b - t + \varepsilon} \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{b + t + \varepsilon}{2} \right) d\tau, \end{aligned}$$

which gives

$$\begin{aligned} & \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau - \frac{2}{b - t + \varepsilon} \int_{t+\varepsilon}^b f(\tau) d\tau + 2 \left( \frac{b - t - \varepsilon}{b - t + \varepsilon} \right) f(t) \\ &= -\frac{2}{b - t + \varepsilon} \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \delta \right) \left( \tau - \frac{b + t + \varepsilon}{2} \right) d\tau, \end{aligned} \quad (2.18)$$

for  $b - t > \varepsilon > 0$ .

If we add (2.16) with (2.18) we deduce the desired equality (2.14).  $\square$

**Theorem 2.3.** Let  $f : (a, b) \rightarrow \mathbb{C}$  be such that for some  $\gamma, \Gamma \in \mathbb{C}$ ,  $\gamma \neq \Gamma$ , we have that  $f \in \bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$ . Then we have the inequality

$$\begin{aligned} & \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left( \frac{b - t}{t - a} \right) \right. \\ & \quad \left. - \frac{2}{\pi} \left( \frac{1}{b - t} \int_t^b f(\tau) d\tau - \frac{1}{t - a} \int_a^t f(\tau) d\tau \right) \right| \\ & \leq \frac{1}{4\pi} |\Gamma - \gamma| (b - a). \end{aligned} \quad (2.19)$$

In particular, we have

$$\begin{aligned} & \left| (Tf) \left( a, b; \frac{a+b}{2} \right) - \frac{4}{\pi} \left( \frac{1}{b-a} \int_{\frac{a+b}{2}}^b f(\tau) d\tau - \frac{1}{b-a} \int_a^{\frac{a+b}{2}} f(\tau) d\tau \right) \right| \\ & \leq \frac{1}{4\pi} |\Gamma - \gamma| (b - a). \end{aligned} \quad (2.20)$$

*Proof.* By using the equality (2.14) for  $\delta = \frac{\gamma+\Gamma}{2}$  and the fact that  $f \in \bar{\Delta}_{(a,b),d}(\gamma, \Gamma)$ , we have for  $\min\{t-a, b-t\} > \varepsilon > 0$  that

$$\begin{aligned}
 & \left| \int_a^{t-\varepsilon} \frac{f(\tau) - f(t)}{\tau - t} d\tau + \int_{t+\varepsilon}^b \frac{f(\tau) - f(t)}{\tau - t} d\tau \right. \\
 & + 2 \left( \frac{1}{t+\varepsilon-a} \int_a^{t-\varepsilon} f(\tau) d\tau - \frac{1}{b-t+\varepsilon} \int_{t+\varepsilon}^b f(\tau) d\tau \right) \\
 & \left. + 2 \left( \frac{b-t-\varepsilon}{b-t+\varepsilon} - \frac{t-\varepsilon-a}{t+\varepsilon-a} \right) f(t) \right| \\
 & \leq \frac{2}{t+\varepsilon-a} \left| \int_a^{t-\varepsilon} \left( \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right) \left( \tau - \frac{a+t-\varepsilon}{2} \right) d\tau \right| \\
 & + \frac{2}{b-t+\varepsilon} \left| \int_{t+\varepsilon}^b \left( \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right) \left( \tau - \frac{b+t+\varepsilon}{2} \right) d\tau \right| \\
 & \leq \frac{2}{t+\varepsilon-a} \int_a^{t-\varepsilon} \left| \left( \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right) \left( \tau - \frac{a+t-\varepsilon}{2} \right) \right| d\tau \\
 & + \frac{2}{b-t+\varepsilon} \int_{t+\varepsilon}^b \left| \left( \frac{f(\tau) - f(t)}{\tau - t} - \frac{\gamma + \Gamma}{2} \right) \left( \tau - \frac{b+t+\varepsilon}{2} \right) \right| d\tau \\
 & \leq \frac{1}{2} |\Gamma - \gamma| \\
 & \times \left[ \frac{2}{t+\varepsilon-a} \int_a^{t-\varepsilon} \left| \tau - \frac{a+t-\varepsilon}{2} \right| d\tau + \frac{2}{b-t+\varepsilon} \int_{t+\varepsilon}^b \left| \tau - \frac{b+t+\varepsilon}{2} \right| d\tau \right] \\
 & = \frac{1}{2} |\Gamma - \gamma| \left[ \frac{2}{t+\varepsilon-a} \frac{(t-\varepsilon-a)^2}{4} + \frac{2}{b-t+\varepsilon} \frac{(b-t-\varepsilon)^2}{4} \right] \\
 & = \frac{1}{4} |\Gamma - \gamma| \left[ \frac{(t-\varepsilon-a)^2}{t+\varepsilon-a} + \frac{(b-t-\varepsilon)^2}{b-t+\varepsilon} \right].
 \end{aligned} \tag{2.21}$$

By taking the limit over  $\varepsilon \rightarrow 0+$  in (2.21) we get

$$\begin{aligned}
 & \left| PV \int_a^b \frac{f(\tau) - f(t)}{\tau - t} d\tau + 2 \left( \frac{1}{t-a} \int_a^t f(\tau) d\tau - \frac{1}{b-t} \int_t^b f(\tau) d\tau \right) \right| \\
 & \leq \frac{1}{4} |\Gamma - \gamma| \left[ \frac{(t-a)^2}{t-a} + \frac{(b-t)^2}{b-t} \right] = \frac{1}{4} |\Gamma - \gamma| (b-a)
 \end{aligned}$$

for  $t \in (a, b)$  and by (2.13) we deduce the desired result (2.19).  $\square$

**Corollary 2.3.** Let  $f : (a, b) \rightarrow \mathbb{R}$  and  $e : (a, b) \rightarrow \mathbb{R}$ ,  $e(t) = t$  such that  $f - me$ ,  $Me - f$  are monotonic nondecreasing on  $(a, b)$ , where  $m < M$  are given real numbers. Then

$$\begin{aligned}
 & \left| (Tf)(a, b; t) - \frac{f(t)}{\pi} \ln \left( \frac{b-t}{t-a} \right) \right. \\
 & \quad \left. - \frac{2}{\pi} \left( \frac{1}{b-t} \int_t^b f(\tau) d\tau - \frac{1}{t-a} \int_a^t f(\tau) d\tau \right) \right| \\
 & \leq \frac{1}{4\pi} (M - m) (b - a) \tag{2.22}
 \end{aligned}$$

for all  $t \in (a, b)$ .

In particular, we have

$$\begin{aligned} & \left| (Tf) \left( a, b; \frac{a+b}{2} \right) - \frac{4}{\pi} \left( \frac{1}{b-a} \int_{\frac{a+b}{2}}^b f(\tau) d\tau - \frac{1}{b-a} \int_a^{\frac{a+b}{2}} f(\tau) d\tau \right) \right| \\ & \leq \frac{1}{4\pi} (M - m) (b - a). \end{aligned} \quad (2.23)$$

**Remark 2.3.** If the function  $f$  is differentiable on  $(a, b)$  and satisfies condition  $m \leq f'(t) \leq M$  for all  $t \in (a, b)$ , then the inequalities (2.22) and (2.23) are valid.

### 3. CONCLUSIONS AND SOME EXAMPLES

In this paper we established some general inequalities for the finite Hilbert transform of complex valued functions for which the divided differences in any two points of the interval are bounded. Further on, we give some simple examples for the exponential function on a finite interval.

If we consider the function  $f(t) = e^t$ ,  $t \in (a, b)$  a real interval, then

$$(Tf)(a, b; t) = \frac{\exp(t)}{\pi} [E_i(b-t) - E_i(a-t)], \quad (3.24)$$

where  $E_i$  is defined by

$$E_i(x) := PV \int_{-\infty}^x \frac{\exp(s)}{s} ds, \quad x \in \mathbb{R}.$$

Indeed, we have

$$\begin{aligned} E_i(b-t) - E_i(a-t) &= PV \int_{a-t}^{b-t} \frac{\exp(s)}{s} ds = PV \int_a^b \frac{\exp(\tau-t)}{\tau-t} ds \\ &= \exp(-t) \pi (T \exp)(a, b; t) \end{aligned}$$

and the equality (3.24) is proved.

We have that  $f'(t) = e^t$ ,  $t \in (a, b)$ , which shows that  $m \leq \exp(a) \leq f'(t) \leq \exp(b) = M$ .

By utilising (1.4) we have

$$\begin{aligned} & \left| E_i(b-t) - E_i(a-t) - \ln \left( \frac{b-t}{t-a} \right) - \frac{\exp(a-t) + \exp(b-t)}{2} (b-a) \right| \\ & \leq \frac{\exp(b-t) - \exp(a-t)}{2} (b-a), \end{aligned} \quad (3.25)$$

while from (2.22) we get

$$\begin{aligned} & \left| E_i(b-t) - E_i(a-t) - \ln \left( \frac{b-t}{t-a} \right) \right. \\ & \quad \left. - 2 \left( \frac{\exp(b-t) - 1}{b-t} - \frac{1 - \exp(a-t)}{t-a} \right) \right| \\ & \leq \frac{1}{4} (\exp(b-t) - \exp(a-t)) (b-a) \end{aligned} \quad (3.26)$$

for  $t \in (a, b)$ .



If we take in (3.25) and (3.26)  $t = \frac{a+b}{2}$ , then we get

$$\begin{aligned} & \left| E_i \left( \frac{b-a}{2} \right) - E_i \left( -\frac{b-a}{2} \right) - \frac{\exp \left( -\frac{b-a}{2} \right) + \exp \left( \frac{b-a}{2} \right)}{2} (b-a) \right| \\ & \leq \frac{1}{2} \left[ \exp \left( \frac{b-a}{2} \right) - \exp \left( -\frac{b-a}{2} \right) \right] (b-a), \end{aligned}$$

and

$$\begin{aligned} & \left| E_i \left( \frac{b-a}{2} \right) - E_i \left( -\frac{b-a}{2} \right) - 4 \left( \frac{\exp \left( \frac{b-a}{2} \right) + \exp \left( -\frac{b-a}{2} \right)}{b-a} \right) \right| \\ & \leq \frac{1}{4} \left[ \exp \left( \frac{b-a}{2} \right) - \exp \left( -\frac{b-a}{2} \right) \right] (b-a), \end{aligned}$$

which, by taking  $x = \frac{b-a}{2} > 0$ , gives

$$|E_i(x) - E_i(-x) - [\exp(-x) + \exp(x)]x| \leq [\exp(x) - \exp(-x)]x \tag{3.27}$$

and

$$\left| E_i(x) - E_i(-x) - 2 \left( \frac{\exp(x) + \exp(-x)}{x} \right) \right| \leq \frac{1}{2} [\exp(x) - \exp(-x)]x \tag{3.28}$$

for  $x > 0$ .

For the function  $f(t) = \frac{1}{t}$ , with  $t \in (a, b) \subset (0, \infty)$  we have

$$(Tf)(a, b; t) = \frac{1}{\pi t} \ln \left( \frac{b-t}{t-a} \right) - \frac{1}{\pi t} \ln \left( \frac{b}{a} \right).$$

Since  $f'(t) = -\frac{1}{t^2}$ , then  $m = -\frac{1}{a^2} \leq f'(t) \leq -\frac{1}{b^2} = M$ , then by (1.4) we have

$$\left| \ln \left( \frac{b}{a} \right) - t \frac{b^2 + a^2}{2a^2b^2} (b-a) \right| \leq t \frac{b+a}{2a^2b^2} (b-a)^2 \tag{3.29}$$

while from (2.22) we get

$$\left| \ln \left( \frac{b}{a} \right) - 2t \left( \frac{\ln t - \ln a}{t-a} - \frac{\ln b - \ln t}{b-t} \right) \right| \leq t \frac{b+a}{4a^2b^2} (b-a)^2 \tag{3.30}$$

for  $t \in (a, b) \subset (0, \infty)$ .

### REFERENCES

- [1] Dragomir, N. M., Dragomir, S. S. and Farrell, P. M., *Some inequalities for the finite Hilbert transform*, Inequality theory and applications, Vol. I, 113–122, Nova Sci. Publ., Huntington, NY, 2001
- [2] Dragomir, N. M., Dragomir, S. S. and Farrell, P. M., *Approximating the finite Hilbert transform via trapezoid type inequalities*, Comput. Math. Appl., **43** (2002), No. 10-11, 1359–1369
- [3] Dragomir, N. M., Dragomir, S. S., Farrell, P. M. and Baxter, G. W., *On some new estimates of the finite Hilbert transform*, Libertas Math., **22** (2002), 65–75
- [4] Dragomir, N. M., Dragomir, S. S., Farrell, P. M. and Baxter, G. W., *A quadrature rule for the finite Hilbert transform via trapezoid type inequalities*, J. Appl. Math. Comput., **13** (2003), No. 1-2, 67–84
- [5] Dragomir, N. M., Dragomir, S. S., Farrell, P. M. and Baxter, G. W., *A quadrature rule for the finite Hilbert transform via midpoint type inequalities*, Fixed point theory and applications, Vol. 5, 11–22, Nova Sci. Publ., Hauppauge, NY, 2004
- [6] Dragomir, S. S., *Inequalities for the Hilbert transform of functions whose derivatives are convex.*, J. Korean Math. Soc., **39** (2002), No. 5, 709–729
- [7] Dragomir, S. S., *Approximating the finite Hilbert transform via an Ostrowski type inequality for functions of bounded variation*, J. Inequal. Pure Appl. Math., **3** (2002), No. 4, Article 51, 19 pp.
- [8] Dragomir, S. S., *Approximating the finite Hilbert transform via Ostrowski type inequalities for absolutely continuous functions*, Bull. Korean Math. Soc., **39** (2002), No. 4, 543–559

- [9] Dragomir, S. S., *Some inequalities for the finite Hilbert transform of a product*, Commun. Korean Math. Soc., **18** (2003), No. 1, 39–57
- [10] Dragomir, S. S. *Sharp error bounds of a quadrature rule with one multiple node for the finite Hilbert transform in some classes of continuous differentiable functions*, Taiwanese J. Math., **9** (2005), No. 1, 95–109
- [11] Dragomir, S. S., *Inequalities and approximations for the Finite Hilbert transform: a survey of recent results*, Preprint RGMIA Res. Rep. Coll., **21** (2018), Art. 30, 90 pp. [<http://rgmia.org/papers/v21/v21a30.pdf>].
- [12] Dragomir, S. S., *Inequalities for the finite Hilbert transform of convex functions*, Preprint RGMIA Res. Rep. Coll., **21** (2018), Art. 31
- [13] Gakhov, F. D., *Boundary Value Problems* (English translation), Pergamon Press, Oxford, 1966
- [14] Liu, W. and Gao, X., *Approximating the finite Hilbert transform via a companion of Ostrowski's inequality for function of bounded variation and applications*, Appl. Math. Comput., **247** (2014), 373–385
- [15] Liu, W., Gao, X. and Wen, Y., *Approximating the finite Hilbert transform via some companions of Ostrowski's inequalities*, Bull. Malays. Math. Sci. Soc., **39** (2016), No. 4, 1499–1513
- [16] Liu, W. and Lu, N., *Approximating the finite Hilbert transform via Simpson type inequalities and applications*. Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys., **77** (2015), No. 3, 107–122
- [17] Mikhlin, S. G. and Prössdorf, S., *Singular Integral Operators* (English translation), Springer Verlag, Berlin, 1986
- [18] Wang, S., Gao, X. and Lu, N., *A quadrature formula in approximating the finite Hilbert transform via perturbed trapezoid type inequalities*, J. Comput. Anal. Appl., **22** (2017), No. 2, 239–246
- [19] Wang, S., Lu, N. and Gao, X., *A quadrature rule for the finite Hilbert transform via Simpson type inequalities and applications*, J. Comput. Anal. Appl., **22** (2017), No. 2, 229–238

VICTORIA UNIVERSITY  
COLLEGE OF ENGINEERING AND SCIENCE  
VICTORIA 8001, MELBOURNE, AUSTRALIA  
Email address: sever.dragomir@vu.edu.au