

# An area formula for the triangle of residual centroids and its generalizations

SZILÁRD ANDRÁS and SÁNDOR NAGYDOBAI KISS

**ABSTRACT.** In this paper we consider an inscribed triangle  $XYZ$  to a triangle  $ABC$  and we establish a relation between the area of these two triangles and the area of the triangle determined by the centroids of the residual triangles  $AZY$ ,  $BXZ$  and  $CYX$ . Moreover we generalize this relation to the case of a general barycenter instead of centroid and also to quadrilaterals.

## 1. INTRODUCTION

In this paper we consider the points  $X, Y, Z$  on the sides  $BC, CA$  and  $AB$  of an arbitrary triangle  $ABC$  and we denote by  $G_A, G_B, G_C$  the centroids of the residual triangles  $AZY, BXZ$  and  $CYX$  respectively. We establish a relation (see Theorem 2) between the area of the triangles  $G_A G_B G_C, XYZ$  and  $ABC$ . This is motivated by some earlier results published by M. Dalcín in [3] and E. Eckart in [5] and [6] concerning triangle centers of residual triangles. More exactly in [3] the author asserted the following property without proof:

**Theorem 1.1** (Proposition 8). *An inscribed triangle and its triangle of residual orthocenters have equal areas.*

This is a consequence of a general property of hexagons having three pairs of parallel sides:

**Theorem 1.2.** *If the pairs of opposite sides of the hexagon  $ABCDEF$  are parallel, then the triangles  $ACE$  and  $BDF$  have equal area.*

This property was a contest problem in 1958 at the famous József Kürschák Mathematical Competition and the solution can be found in [2].

In this paper we establish a relation between the area of the initial triangle  $ABC$ , the inscribed triangle  $XYZ$  and the triangle determined by the centroids of the residual triangles  $AZY, BXZ$  and  $CYX$ . Moreover we give a possible generalization for quadrilaterals.

## 2. MAIN RESULTS

**Theorem 2.3.** *The area of the triangle of residual centroids satisfies the following relation:*

$$9 \cdot \text{Area}(G_A G_B G_C) = 2 \cdot \text{Area}(ABC) + \text{Area}(XYZ). \quad (2.1)$$

---

Received: 20.09.2020. In revised form: 21.01.2021. Accepted: 28.01.2021

2010 *Mathematics Subject Classification.* 51M04, 51M25.

Key words and phrases. *residual triangles, centroids, area.*

Corresponding author: Szilárd András; [andrasz@math.ubbcluj.ro](mailto:andrasz@math.ubbcluj.ro)




FIGURE 1. The centroids of the residual triangles

*Proof.* Let  $x, y, z \in \mathbb{R}$  such that the normalized barycentric coordinates of the points  $X, Y, Z$  are

$$X = (0, 1 - x, x), Y = (y, 0, 1 - y), Z = (1 - z, z, 0).$$

With these notations we have

$$\begin{aligned} \text{Area}(XYZ) &= \text{Area}(ABC) \cdot \begin{vmatrix} 0 & 1 - x & x \\ y & 0 & 1 - y \\ 1 - z & z & 0 \end{vmatrix} \\ &= \text{Area}(ABC) \cdot (xyz + (1 - x)(1 - y)(1 - z)). \end{aligned} \quad (2.2)$$

The barycentric coordinates of the centroid  $G_A, G_B$  and  $G_C$  can be expressed as follows:

$$G_A = \frac{1}{3}(2 + y - z, z, 1 - y), G_B = \frac{1}{3}(1 - z, 2 + z - x, x), G_C = \frac{1}{3}(y, 1 - x, 2 + x - y)$$

so the area of the triangle  $G_A G_B G_C$  is

$$\begin{aligned} \text{Area}(G_A G_B G_C) &= \frac{1}{27} \text{Area}(ABC) \cdot \begin{vmatrix} 2 + y - z & z & 1 - y \\ 1 - z & 2 + z - x & x \\ y & 1 - x & 2 + x - y \end{vmatrix} \\ &= \frac{1}{9} \text{Area}(ABC) \cdot \begin{vmatrix} 1 & z & 1 - y \\ 1 & 2 + z - x & x \\ 1 & 1 - x & 2 + x - y \end{vmatrix} \\ &= \frac{1}{9} \text{Area}(ABC) \cdot (2 + xyz + (1 - x)(1 - y)(1 - z)). \end{aligned} \quad (2.3)$$

From (2.3) and (2.2) we obtain (2.1), so the proof is complete.  $\square$

**Remark 2.1.** The area is considered as oriented one, so the relation remains true even if, the points  $X, Y, Z$  are outside the segments  $BC, CA, AB$ .

**Remark 2.2.** Using Theorem 1.1. we obtain the following equivalent formulation:

$$9 \cdot \text{Area}(G_A G_B G_C) = 2 \cdot \text{Area}(ABC) + \text{Area}(H_A H_B H_C), \quad (2.4)$$

where  $H_A, H_B$  and  $H_C$  are the orthocenters of the residual triangles  $AZY, BXZ$  and  $CYX$  respectively.

**Theorem 2.4.** Consider the arbitrary points  $X, Y, Z$  on the sides  $BC, CA$  and  $AB$  of an arbitrary triangle  $ABC$ . The points  $A_1, A_2$  and  $A_3$  have normalized barycentric coordinates  $(\lambda_1, \lambda_2, \lambda_3)$ ,  $(\lambda_3, \lambda_1, \lambda_2)$  and  $(\lambda_2, \lambda_3, \lambda_1)$  respectively with respect to the triangle  $AZY$ . In a similar way we




FIGURE 2. Replacing the centroid with 3 points

consider the points  $B_1, B_2$  and  $B_3$  in the triangle  $BXZ$  and  $C_1, C_2, C_3$  in the triangle  $CYX$ . With these notations we have

$$\text{Area}(A_1B_1C_1) + \text{Area}(A_2B_2C_2) + \text{Area}(A_3B_3C_3) = c_1 \cdot \text{Area}(ABC) + c_2 \cdot \text{Area}(XYZ), \quad (2.5)$$

where

$$\begin{aligned} c_1 &= \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_1 \cdot \lambda_2 + \lambda_2 \cdot \lambda_3 + \lambda_3 \cdot \lambda_1, \\ c_2 &= 2(\lambda_1^2 + \lambda_2^2 + \lambda_3^2) - (\lambda_1 \cdot \lambda_2 + \lambda_2 \cdot \lambda_3 + \lambda_3 \cdot \lambda_1). \end{aligned}$$

**Remark 2.3.** If  $\lambda_1 = \lambda_2 = \lambda_3 = \frac{1}{3}$ , we have  $A_1 = A_2 = A_3 = G_A$  the centroid of  $AYZ$ ,  $B_1 = B_2 = B_3 = G_B$  the centroid of  $BZX$  and  $C_1 = C_2 = C_3 = G_C$  the centroid of  $CXY$ . Moreover in this case  $c_1 = \frac{2}{3}$  and  $c_2 = \frac{1}{3}$ , so we obtain (2.1).

*Proof of theorem 2.3.* Using the same notations as in proof of Theorem 2.3 we obtain for the coordinates of the points  $A_1, B_1$  and  $C_1$  the following expressions:

$$\begin{aligned} A_1 &= (\lambda_1 + \lambda_2 \cdot (1 - z) + \lambda_3 \cdot y, \lambda_2 \cdot z, \lambda_3 \cdot (1 - y)), \\ B_1 &= (\lambda_1 \cdot (1 - z), \lambda_1 + \lambda_2 \cdot (1 - x) + \lambda_3 \cdot z, \lambda_2 \cdot x), \\ C_1 &= (\lambda_1 \cdot y, \lambda_2 \cdot (1 - x), \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot x). \end{aligned}$$

So for the ratio  $R_1 = \frac{\text{Area}(A_1B_1C_1)}{\text{Area}(ABC)}$  we obtain (using  $\lambda_1 + \lambda_2 + \lambda_3 = 1$ ):

$$\begin{aligned} R_1 &= \begin{vmatrix} \lambda_1 + \lambda_2 \cdot (1 - z) + \lambda_3 \cdot y & \lambda_2 \cdot z & \lambda_3 \cdot (1 - y) \\ \lambda_1 \cdot (1 - z) & \lambda_1 + \lambda_2 \cdot (1 - x) + \lambda_3 \cdot z & \lambda_2 \cdot x \\ \lambda_1 \cdot y & \lambda_2 \cdot (1 - x) & \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot x \end{vmatrix} \\ &= \begin{vmatrix} 1 & \lambda_2 \cdot z & \lambda_3 \cdot (1 - y) \\ 1 & \lambda_1 + \lambda_2 \cdot (1 - x) + \lambda_3 \cdot z & \lambda_2 \cdot x \\ 1 & \lambda_2 \cdot (1 - x) & \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot x \end{vmatrix} \\ &= \begin{vmatrix} \lambda_1 + \lambda_2 \cdot (1 - x - z) + \lambda_3 \cdot z & \lambda_2 \cdot x - \lambda_3 \cdot (1 - y) \\ \lambda_2 \cdot (1 - x - z) & \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot (x + y - 1) \end{vmatrix} \\ &= (\lambda_1 + \lambda_3 \cdot z) \begin{vmatrix} 1 & \lambda_2 \cdot x - \lambda_3 \cdot (1 - y) \\ 0 & \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot (x + y - 1) \end{vmatrix} + \\ &\quad \lambda_2(1 - x - z) \begin{vmatrix} 1 & \lambda_2 \cdot x - \lambda_3 \cdot (1 - y) \\ 1 & \lambda_1 + \lambda_2 \cdot (1 - y) + \lambda_3 \cdot (x + y - 1) \end{vmatrix}. \end{aligned} \quad (2.6)$$

So we have

$$R_1 = \lambda_1^2 + \lambda_2^2(1-x-y)(1-x-z) + \lambda_3^2(zx+zy-z) + \lambda_1 \cdot \lambda_2(2-x-y-z) + \lambda_2 \cdot \lambda_3(z+x-zy-xz-x^2) + \lambda_3 \cdot \lambda_1(x+y+z-1).$$

By a similar calculation we deduce a relation for  $R_2 = \frac{\text{Area}(A_2B_2C_2)}{\text{Area}(ABC)}$  and for  $R_3 = \frac{\text{Area}(A_3B_3C_3)}{\text{Area}(ABC)}$ . Due to the symmetry of the notations by adding the three ratios we obtain

$$R_1 + R_2 + R_3 = c_1 + c_2(1-x-y-z+xy+yz+zx),$$

which completes the proof.  $\square$

**Example 2.1.** If  $M_1$  is the midpoint of  $ZY$ , the midpoint of  $AM_1$  has barycentric coordinates  $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ , so applying the previous result we deduce

$$\text{Area}(A_1B_1C_1) + \text{Area}(A_2B_2C_2) + \text{Area}(A_3B_3C_3) = \frac{11}{16} \cdot \text{Area}(ABC) + \frac{7}{16} \cdot \text{Area}(XYZ), \quad (2.7)$$

where  $A_1, A_2, A_3, B_1, B_2, B_3$  and  $C_1, C_2, C_3$  are the midpoints of the corresponding medians in the triangles  $AZY, BXZ$  and  $CYX$  respectively.

In what follows we give a more general version for the relation (2.1).

**Theorem 2.5.** If  $G_1, G_2$  and  $G_3$  are the centroids of the triangles  $A_{11}A_{12}A_{13}, A_{21}A_{22}A_{23}$  and  $A_{31}A_{32}A_{33}$ , then

$$9\text{Area}[G_1G_2G_3] = \text{Area}[A_{11}A_{21}A_{31}] + \text{Area}[A_{12}A_{22}A_{32}] + \text{Area}[A_{13}A_{23}A_{33}] + \text{Area}[A_{11}A_{22}A_{33}] + \text{Area}[A_{12}A_{23}A_{31}] + \text{Area}[A_{13}A_{21}A_{32}] + \text{Area}[A_{11}A_{23}A_{32}] + \text{Area}[A_{12}A_{21}A_{33}] + \text{Area}[A_{13}A_{22}A_{31}] \quad (2.8)$$

*Proof.* For the area of an arbitrary triangle  $UVT$  we use its expression in complex numbers (see [1], page 109, [7], [4]):

$$\text{Area}[UVT] = \frac{1}{2} \text{Im}(u\bar{v} + v\bar{t} + t\bar{u}),$$

where  $u, v, t$  are the affixes of the points  $U, V, T$ . Using this expression and the affixes of the centroids, we have

$$9\text{Area}[G_1G_2G_3] = \frac{1}{2} \text{Im}((a_{11} + a_{12} + a_{13})\overline{(a_{21} + a_{22} + a_{23})} + (a_{21} + a_{22} + a_{23})\overline{(a_{31} + a_{32} + a_{33})} + (a_{31} + a_{32} + a_{33})\overline{(a_{11} + a_{12} + a_{13})}).$$

So

$$\begin{aligned} 9\text{Area}[G_1G_2G_3] &= \frac{1}{2} \text{Im} \sum_{i=1}^3 \left( \sum_{j=1}^3 a_{ij} \right) \left( \sum_{j=1}^3 \overline{a_{i+1j}} \right) \\ &= \frac{1}{2} \text{Im} ((a_{11}\overline{a_{21}} + a_{21}\overline{a_{31}} + a_{31}\overline{a_{11}}) + (a_{12}\overline{a_{22}} + a_{22}\overline{a_{32}} + a_{32}\overline{a_{12}}) \\ &\quad + (a_{13}\overline{a_{23}} + a_{23}\overline{a_{33}} + a_{33}\overline{a_{13}})) \\ &+ \frac{1}{2} \text{Im} ((a_{11}\overline{a_{22}} + a_{22}\overline{a_{33}} + a_{33}\overline{a_{11}}) + (a_{12}\overline{a_{23}} + a_{23}\overline{a_{31}} + a_{31}\overline{a_{12}}) \\ &\quad + (a_{13}\overline{a_{21}} + a_{21}\overline{a_{32}} + a_{32}\overline{a_{13}})) + \\ &+ \frac{1}{2} \text{Im} ((a_{11}\overline{a_{23}} + a_{23}\overline{a_{32}} + a_{32}\overline{a_{11}}) + (a_{12}\overline{a_{21}} + a_{21}\overline{a_{33}} + a_{33}\overline{a_{12}}) \\ &\quad + (a_{13}\overline{a_{22}} + a_{22}\overline{a_{31}} + a_{31}\overline{a_{13}})). \end{aligned}$$




FIGURE 3. Special configuration of three triangles

Identifying the terms in the right-hand side as areas we obtain (2.8), so the proof is complete.  $\square$

**Remark 2.4.** Using the notations of the figure 3 ( $X = A_{22} = A_{33}, \dots$ ) we obtain

$$9\text{Area}[G_1G_2G_3] = \text{Area}[ABC] + 2\text{Area}[XYZ] + \text{Area}[AZY] + \text{Area}[BXZ] + \text{Area}[CYX], \quad (2.9)$$

so if the points  $XYZ$  are on the sides of the triangle  $ABC$ , we have

$$\text{Area}[ABC] = \text{Area}[XYZ] + \text{Area}[AZY] + \text{Area}[BXZ] + \text{Area}[CYX],$$

which compared to (2.9) implies (2.1).

On the other hand the relation (2.9) is valid even in the case when  $X \notin BC, Y \notin CA$  and  $Z \notin AB$ , so it implies the following interesting property for a hexagon  $AZBXCXY$  :

**Corollary 2.1.** *If  $AZBXCXY$  is an arbitrary hexagon,  $G_1, G_2$  and  $G_3$  are the centroids of the triangles  $AZY, BXZ$  and  $CYX$  respectively, then relation (2.9) holds.*

Applying this corollary twice and interchanging the role of the points  $ABC$  and  $XYZ$  we obtain the following property:

**Corollary 2.2.** *If in the hexagon  $ABCDEF$   $G_1, G_2, G_3, G_4, G_5$  and  $G_6$  are the centroids of the triangles  $ABF, BCA, CDB, DEC, EFD$  and  $FAE$  respectively (see figure 4), then*

$$\text{Area}(G_1G_3G_5) = \text{Area}(G_2G_4G_6).$$

This property also reduces to Theorem 1.2. since the opposite sides of the hexagon  $G_1G_2G_3G_4G_5G_6$  are parallel to the diagonals  $AD, BE$  and  $CF$ .

Theorem 2.5. can also be generalized to more than 3 triangles, but the number of triangles in the right-hand side expression is too large (16). For this reason we do not assert this case or the higher dimensional analogous properties, we prove only a property for quadrilaterals that is analogous to Theorem 2.3.

**Theorem 2.6.** *If  $ABCD$  is a quadrilateral,  $M \in BC, N \in CD, P \in DA, Q \in AB$  and we denote by  $G_A, G_B, G_C, G_D$  the centroids of the triangles  $AQP, BMQ, CNM$  and  $DPN$ , then*

$$9\text{Area}[G_A G_B G_C G_D] = 2\text{Area}[ABCD] + 2\text{Area}[MNPQ] + \text{Area}[AMDQCPBNA] \quad (2.10)$$





FIGURE 4. An interesting property of the hexagon



**Remark 2.5.** For a better understanding of the term  $Area[AMDQCPBNA]$  from the previous theorem we illustrate it in Figure 5. In fact it is the sum of the areas  $A_1, A_2$  and  $A_3$  shown in this figure.

*Proof.* Using complex numbers as in the proof of theorem 2.5. we have




FIGURE 5.  $Area(AMDQCPBNA) = A_1 + A_2 + A_4$

$$9Area(G_A G_B G_C G_D) = \frac{9}{2} Im(g_A \cdot \bar{g}_B + g_B \cdot \bar{g}_C + g_C \cdot \bar{g}_D + g_D \cdot \bar{g}_A) \quad (2.11)$$

$$= \frac{1}{2} Im((p + a + q) \cdot \overline{q + b + m} + (q + b + m) \cdot \overline{m + c + n}) \quad (2.12)$$

$$+ \frac{1}{2} Im((m + c + n) \cdot \overline{n + d + p} + (n + d + p) \cdot \overline{p + a + q}) \quad (2.13)$$

Expanding the products in the right-hand side we obtain 36 terms. We can omit the terms  $q \cdot \bar{q}$ ,  $m \cdot \bar{m}$ ,  $n \cdot \bar{n}$  and  $p \cdot \bar{p}$  since these terms are real numbers, so their imaginary part is 0. The sum of terms containing  $p \cdot \bar{q}$ ,  $q \cdot \bar{m}$ ,  $m \cdot \bar{n}$  and  $n \cdot \bar{p}$  is  $Area(MNPQ)$ , while the sum of terms containing  $a \cdot \bar{b}$ ,  $b \cdot \bar{c}$ ,  $c \cdot \bar{d}$  and  $d \cdot \bar{a}$  is  $Area(ABCD)$ . We have also the terms  $q \cdot \bar{m}$ ,  $m \cdot \bar{n}$ ,  $n \cdot \bar{p}$  and  $p \cdot \bar{q}$ , whose sum gives the second  $Area(MNPQ)$  on the right-hand side.

In order to identify the second  $Area(ABCD)$  observe that the terms  $a \cdot \bar{q}$ ,  $q \cdot \bar{n}$ ,  $n \cdot \bar{d}$ ,  $d \cdot \bar{p}$  and  $p \cdot \bar{a}$  form  $Area(AQN DPA)$ , while the terms  $q \cdot \bar{b}$ ,  $b \cdot \bar{m}$ ,  $m \cdot \bar{c}$ ,  $c \cdot \bar{n}$  and  $n \cdot \bar{q}$  form  $Area(QBMCN Q)$ , so the sum of these two areas is  $Area(ABCD)$ . The remaining terms are  $a \cdot \bar{m}$ ,  $m \cdot \bar{d}$ ,  $d \cdot \bar{q}$ ,  $q \cdot \bar{c}$ ,  $c \cdot \bar{p}$ ,  $p \cdot \bar{b}$ ,  $b \cdot \bar{n}$  and  $n \cdot \bar{a}$  whose sum gives  $Area(AMDQCPBNA)$ , so the proof is complete.  $\square$

In the previous proof we used that the points  $M, N, P, Q$  are on the sides of the quadrilateral only in identifying the area of  $AQNDPA$  with the area of  $AQND$ , so a more general identity is valid for any octagon  $AQBMCNDP$ .

**Theorem 2.7.** *If in the octagon  $AQBMCNDP$   $G_A, G_B, G_C$  and  $G_D$  are the centroids of the triangles  $AQP, BMQ, CNM$  and  $DPN$  respectively, then*

$$9\text{Area}[G_A G_B G_C G_D] = \text{Area}[ABCD] + \text{Area}[AQBMCNDP] \\ + 2\text{Area}[MNPQ] + \text{Area}[AMDQCPBNA]$$

Applying this theorem twice we obtain the following property:

**Corollary 2.3.** *If in the octagon  $A_1 A_2 A_3 A_4 A_5 A_6 A_7 A_8$  we denote for all  $i \in \{1, 2, 3, \dots, 8\}$  by  $G_i$  the centroid of the triangle  $A_{i-1} A_i A_{i+1}$  (the indices are taken circularly, so  $A_9 = A_1$  and  $A_0 = A_8$ ), then*

$$\text{Area}(A_1 A_3 A_5 A_7) - \text{Area}(A_2 A_4 A_6 A_8) = 9(\text{Area}(G_2 G_4 G_6 G_8) - \text{Area}(G_1 G_3 G_5 G_7)).$$

#### REFERENCES

- [1] Andreescu, T. and Andrica, D., *Complex numbers from A to ... Z*, Second Edition, Birkhauser, 2014
- [2] Barrington Leigh, R. and Liu, A., *Hungarian problem book IV*, MAA Press, 2011 (or Kömal Journal, 1959, page 70–77)
- [3] Dalcín, M., *Isotomic Inscribed Triangles and Their Residuals*, Forum Geometricorum, **3** (2003), 125–134
- [4] Deaux, R., *Introduction to the Geometry of Complex Numbers*, Ungar, New York, 1956
- [5] Schmidt, E., *Circumcenters of Residual Triangles*, Forum Geometricorum, **3** (2003), 207–214
- [6] Schmidt, E., *Miquel points and inscribed triangles*, <http://eckartschmidt.de/Miquel.pdf>, Accessed 20.09.2020
- [7] Tonov, I., *Complex numbers* (in bulgarian), Ed. Narodna prosveta, Sofia, 1979

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE  
 BABEŞ-BOLYAI UNIVERSITY  
 M. KOGĂLNICEANU, NO 1, 400084, CLUJ-NAPOCA, ROMANIA  
 E-mail address: andrasz@math.ubbcluj.ro

SATU MARE, ROMANIA  
 E-mail address: d.sandor.kiss@gmail.com